

A STRUCTURED INVERSE SPECTRUM PROBLEM FOR INFINITE GRAPHS

KEIVAN HASSANI MONFARED AND EHSSAN KHANMOHAMMADI

ABSTRACT. In this paper it is shown that for a given infinite graph G on countably many vertices, and a compact, infinite set Λ of real numbers there is a real symmetric matrix A whose graph is G and its spectrum is Λ . Moreover, the set of limit points of Λ equals the essential spectrum of A , and the isolated points of Λ are eigenvalues of A with multiplicity one. It is also shown that any two such matrices constructed by our method are approximately unitarily equivalent.

Keywords: Inverse Spectrum Problem, Jacobian Method, Infinite Graph, Isospectral, Essential Spectrum, Approximate Unitary Equivalence.

AMS MSC: 05C50, 15A18, 15A29, 47A10.

1. INTRODUCTION

Inverse eigenvalue problems stemming from applications have been studied to a great extent from several points of view. For instance, the inverse Sturm-Liouville problem essentially seeks to determine the density of a vibrating string from its natural frequencies (see [3, p. 83], and [12]). Similar but more complicated problems occur in other areas of science and engineering (see for example [1] for problems arising from geophysics, and [13] for an example in applied electromagnetics). Common approaches involve solving a finite discretization of the problem and proving that the solution of the discrete problem converges to the solution of the continuous problem. For example, Hald in his 1978 paper [8] used the Rayleigh-Ritz method for calculating the eigenvalues of a two point boundary value problem, and reduced the inverse problem for the differential equation to a discrete inverse eigenvalue problem.

It is of interest to analyze the vibrations of lumped parameter systems, that is, systems that are modelled as rigid masses joined by massless springs. In general, more accurate results are obtained by increasing the number of masses and springs, that is, by increasing

the degrees of freedom. As the number of degrees of freedom is increased without limit, the concept of the system with distributed mass and elasticity is formed. Stokey [16] gives a thorough treatment of the direct problem when the degrees of freedom is infinite.

The λ -structured-inverse-eigenvalue-problem (λ -SIEP) asks about the existence of a matrix whose graph and eigenvalues are given. In a lot of cases, one seeks to find a symmetric tridiagonal matrix (a matrix whose graph is a disjoint union of paths) with prescribed eigenvalues [5]. In 1989 Duarte [7] showed that the λ -SIEP problem has a solution whenever the given graph is a finite tree. In [15] the general problem is described and solved when all the eigenvalues are distinct and the graph is finite.

In this paper our goal is to establish analogous results when the graph and hence the solution matrix are infinite. In order to do so, in Section 2 we introduce a property that captures a notion of *genericity* for finite matrices in the settings of the Jacobian method [15]. Then in Section 3 we solve the finite λ -SIEP again but instead of solving the problem in one step, we find the solutions A_n using induction on n , the number of vertices of the graph, and the Jacobian method. This approach enables us to control the norm of A_n 's in each step. Finally, in Section 4 we will show that the limit of A_n 's as n approaches infinity exists and has the given graph and spectrum.

Throughout the paper \circ denotes the Schur (entry-wise) product of two matrices, and the Lie bracket $[A, B]$ is the commutator of two matrices A and B , that is $AB - BA$. A zero matrix of appropriate size is denoted by O , all vectors are written in bold small letters, and $\mathbf{0}$ denotes a zero vector of appropriate size to the context.

We assume throughout this paper that all matrices have real entries. Let A be an $n \times n$ symmetric matrix and G a simple graph on n vertices. For a set $S \subset [n] = \{1, 2, \dots, n\}$ let $A[S]$ denote the submatrix of A obtained by deleting the rows and columns indexed by $[n] \setminus S$. Also let $G[S]$ denote the induced subgraph of G on the vertices labeled by S . We say G is the graph of A when for $i \neq j$ we have $A_{ij} \neq 0$ if and only if vertices i and j are adjacent in G . Furthermore, for an infinite matrix A whose rows and columns are indexed by \mathbb{N} , we say that an infinite graph G on countably many vertices labeled by \mathbb{N} is the graph of A when $G[S]$ is the graph of $A[S]$ for any nonempty finite $S \subset \mathbb{N}$. Note that G is not necessarily locally finite.

2. THE WEAK SPECTRAL PROPERTY

The problem studied in this paper is about the existence of a matrix with a given spectrum and the property that each off-diagonal entry of the matrix is prescribed to be either zero or nonzero with no restriction on the diagonal entries. The Jacobian method [15] starts with a simple matrix with the given spectrum, and changes the entries slightly to obtain the desired pattern of zero and nonzero entries while keeping the spectrum intact. In this application of the Jacobian method we start with a diagonal matrix with the given spectrum, and change the off-diagonal entries one row and one column at a time, and adjust the diagonal entries. The process of showing that changing off-diagonal entries can be compensated with a change in the diagonal entries so that the spectrum remains intact involves using the Implicit Function Theorem (IFT). Checking the necessary conditions of the IFT for this problem involves a notion of robustness and genericity for the solution of the inverse problem we are interested in. A stronger version of this robustness is introduced in [2] as the Strong Spectral Property (SSP). In this section we introduce a property for finite matrices, similar to the SSP, which we call the Weak Spectral Property (WSP), and study some of its properties that are critical to our approach in solving the inverse problem. The WSP implies the SSP, because the WSP uses weaker assumptions than the SSP.

Definition 2.1. A symmetric matrix A is said to have the *Weak Spectral Property* (or A has the WSP for short) if $X = O$ is the only symmetric matrix satisfying

- (1) $X \circ I = O$, and
- (2) $[X, A] = O$.

Note that any 1×1 matrix has the WSP. When $n \geq 2$, an $n \times n$ scalar matrix cannot have the WSP, since any such matrix belongs to the center of the algebra of matrices. More generally, any matrix A with the WSP cannot have a constant diagonal, because if all the diagonal entries of A are equal to a constant c , then

$$(A - cI) \circ I = O \text{ and } [A - cI, A] = O.$$

Thus either $X = A - cI$ is a nonzero solution of (1) and (2), or A equals the scalar matrix cI . For 2×2 matrices this necessary condition for the WSP is sufficient as well, and using this it is easy to give explicit examples to show that the WSP is not necessarily invariant under a change of basis.

Observation 2.2. A 2×2 symmetric matrix has the WSP if and only if it has distinct diagonal entries.

However for $n \times n$ matrices with $n \geq 2$ having distinct diagonal entries does not guarantee the WSP, as the following example illustrates.

Example 2.3. Consider the matrices

$$A = \begin{bmatrix} 4 & 0 & 1 \\ 0 & 3 & 0 \\ 1 & 0 & 2 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 4 & 1 & 0 \\ 1 & 3 & 1 \\ 0 & 1 & 2 \end{bmatrix}.$$

Then a short calculation shows that the matrix A has the WSP but not the matrix B .

The following lemma shows that the WSP is an *open property*. In fact, the proof shows that *any* sufficiently small perturbation, symmetric or not, of a matrix with the WSP satisfies the conditions of Definition 2.1.

Lemma 2.4 (The WSP is an open property). *If a matrix A has the WSP, then any sufficiently small symmetric perturbation of A also has the WSP.*

Proof. We prove the contrapositive of the statement. Let $\|\cdot\|$ denote any matrix norm and assume that for each $n \in \mathbb{N}$ there exist nonzero symmetric matrices E_n and X_n such that

- (1) $X_n \circ I = O$,
- (2) $[X_n, A + E_n] = O$, and
- (3) $\|E_n\| = 1/n$.

With a scaling we can set $\|X_n\| = 1$ for all values of n , and X_n still would satisfy the properties (1) and (2). Note that the set of matrices of norm one is compact in the set of all square matrices with the Euclidean topology. Hence we can substitute the sequence $\{X_n\}$, if necessary, with a convergent subsequence that we again denote by $\{X_n\}$. Thus, taking the limit of the commutation relation (2) as $n \rightarrow \infty$, we obtain

$$[\lim_{n \rightarrow \infty} X_n, A] = \lim_{n \rightarrow \infty} [X_n, A] = \lim_{n \rightarrow \infty} [E_n, X_n] = O.$$

This proves that A does not have the WSP, because the nonzero matrix $X = \lim_{n \rightarrow \infty} X_n$ satisfies the conditions of Definition 2.1. \square

Observation 2.5. Note that a matrix A has the WSP if and only if $A + cI$ has the WSP for any real number c . This implies that the set of symmetric matrices without the WSP has no isolated points.

We conclude this section by proving that the direct sum of two matrices with the WSP which do not share an eigenvalue has the WSP.

Lemma 2.6. *Let A and B be two matrices with the WSP which do not have a common eigenvalue. Then $A \oplus B$ also has the WSP.*

Proof. Let

$$X = \left[\begin{array}{c|c} X_1 & X_2 \\ \hline X_3 & X_4 \end{array} \right],$$

where X_1 and X_4 have the same size as A and B , respectively, and assume that $X \circ I = O$, that is, $X_1 \circ I = O$ and $X_4 \circ I = O$. Furthermore, assume that $[X, A \oplus B] = O$. We want to show that $X = O$. Note that

$$[X, A \oplus B] = \left[\begin{array}{c|c} [X_1, A] & X_2B - AX_2 \\ \hline X_3A - BX_3 & [X_4, B] \end{array} \right] = O.$$

Hence $[X_1, A] = O$ and $[X_4, B] = O$. Since A and B have the WSP, we conclude that $X_1 = O$ and $X_4 = O$. Furthermore, $X_2B - AX_2 = O$ and $X_3A - BX_3 = O$. That is, X_2 and X_3 are the intertwining matrices of A and B . By Lemma 1.1 of [15] $X_2 = O$ and $X_3 = O$, since A and B do not have a common eigenvalue. \square

We will use the following corollary of the above lemma in the next section.

Corollary 2.7. *If a matrix A has the WSP, then so does $A \oplus [c]$ where c is any real number that is not an eigenvalue of A .*

3. FINITE λ -SIEP SOLVED WITH INDUCTION

In this section, to control the norm of matrices constructed, we provide an inductive proof for Theorem 3.2 below, which is also proved using the Jacobian method in [15].

Let G be a given graph on n vertices $1, 2, \dots, n$ and with m edges $\{i_\ell, j_\ell\}$ where $\ell = 1, 2, \dots, m$, and $i_\ell < j_\ell$. Also, let $\lambda_1 < \lambda_2 < \dots < \lambda_n$ be n distinct real numbers, and let $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and $\mathbf{y} = (y_1, y_2, \dots, y_m)$, where x_i 's and y_i 's are independent real unknowns. Define $M = M(\mathbf{x}, \mathbf{y})$ to be the $n \times n$ symmetric matrix whose i -th diagonal entry is x_i , its (i_ℓ, j_ℓ) and (j_ℓ, i_ℓ) entries are y_ℓ , and it is zero elsewhere. Define

$$f: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$$

$$(\mathbf{x}, \mathbf{y}) \mapsto (\lambda_1(M), \lambda_2(M), \dots, \lambda_n(M)),$$

where $\lambda_i(M)$ is the i -th smallest eigenvalue of M . We want to show that there is a real symmetric matrix A whose graph is G and its eigenvalues are $\lambda_1, \lambda_2, \dots, \lambda_n$. In other words, we want to find $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^n \times \mathbb{R}^m$

such that no entry of \mathbf{b} is zero and $f(\mathbf{a}, \mathbf{b}) = (\lambda_1, \lambda_2, \dots, \lambda_n)$. In order to do so we introduce a new function

$$g: \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$$

$$(\mathbf{a}, \mathbf{b}) \mapsto \left(\operatorname{tr} M, \frac{1}{2} \operatorname{tr} M^2, \dots, \frac{1}{n} \operatorname{tr} M^n \right).$$

Note that by Newton's identities [14] there exists an invertible function $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $h \circ g(\mathbf{a}, \mathbf{b}) = f(\mathbf{a}, \mathbf{b})$ for all $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^n \times \mathbb{R}^m$. Thus it suffices to show that there is $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that none of the entries of \mathbf{b} is zero, and

$$(3.1) \quad g(\mathbf{a}, \mathbf{b}) = \left(\sum_{i=1}^n \lambda_i, \frac{1}{2} \sum_{i=1}^n \lambda_i^2, \dots, \frac{1}{n} \sum_{i=1}^n \lambda_i^n \right).$$

Let $\operatorname{Jac}_x(g)$ denote the matrix obtained from the Jacobian matrix of g restricted to the columns corresponding to the derivatives with respect to x_i 's. Let $A = M(\mathbf{a}, \mathbf{b})$ for some $(\mathbf{a}, \mathbf{b}) \in \mathbb{R}^n \times \mathbb{R}^m$. The matrix $\operatorname{Jac}_x(g) \Big|_A$ is the evaluation of $\operatorname{Jac}_x(g)$ at (\mathbf{a}, \mathbf{b}) . Then by Lemma 3.1 of [15] it is easy to see that

$$\operatorname{Jac}_x(g) \Big|_A = \begin{bmatrix} I_{11} & \cdots & I_{n-1, n-1} & I_{nn} \\ A_{11} & \cdots & A_{n-1, n-1} & A_{nn} \\ \vdots & \ddots & \vdots & \vdots \\ A_{11}^{n-1} & \cdots & A_{n-1, n-1}^{n-1} & A_{nn}^{n-1} \end{bmatrix},$$

where A_{ij}^k denotes the (i, j) entry of A^k .

Lemma 3.1. *Let A be an $n \times n$ symmetric matrix with n distinct eigenvalues. Then A has the WSP if and only if $\operatorname{Jac}_x(g) \Big|_A$ is invertible.*

Proof. First suppose that A has the WSP. We show that $J = \operatorname{Jac}_x(g) \Big|_A$ is invertible by proving that the rows of J are linearly independent. Let $\boldsymbol{\alpha} = (\alpha_0, \alpha_1, \dots, \alpha_{n-1})$. For $p(x) = \alpha_0 + \alpha_1 x + \cdots + \alpha_{n-1} x^{n-1}$ and $X := p(A)$, it is clear that $[X, A] = O$. Assume that $\boldsymbol{\alpha} J = \mathbf{0}$ and observe that this holds if and only if $X \circ I = O$. Since A has the WSP, this implies that $X = p(A) = O$. Using the fact that A has n distinct eigenvalues and that its minimal polynomial has degree n we conclude that $p(x) \equiv 0$, and hence $\boldsymbol{\alpha} = \mathbf{0}$ as we wanted.

Next, assume that $J = \operatorname{Jac}_x(g) \Big|_A$ is invertible and that $X \circ I = O$ and $[X, A] = O$ for some symmetric matrix X . Since X commutes with the matrix A with n distinct eigenvalues, X must be a polynomial

in A of degree at most $n - 1$ (see for instance [9, Problem 1.3.P4]). Suppose $X = p(A)$ for $p(x) = \alpha_0 + \alpha_1 x + \cdots + \alpha_{n-1} x^{n-1}$. Let $\boldsymbol{\alpha} = (\alpha_0, \alpha_1, \dots, \alpha_{n-1})$. By the observation in the previous part, $X \circ I = O$ implies $\boldsymbol{\alpha} J = \mathbf{0}$. But since J is invertible, $\boldsymbol{\alpha} = \mathbf{0}$, and hence $p(x) \equiv 0$. Therefore, $X = p(A) = O$. This proves that A has the WSP. \square

Theorem 3.2. *Given a graph G on n vertices and a set of n distinct real numbers $\Lambda = \{\lambda_1, \lambda_2, \dots, \lambda_n\}$, there is an $n \times n$ symmetric matrix whose graph is G and its spectrum is Λ .*

Proof. By induction on the number of vertices we prove the stronger result that there exist \mathbf{a} and \mathbf{b} satisfying (3.1) such that the graph of $A = M(\mathbf{a}, \mathbf{b})$ is G and A has the WSP.

When $n = 1$, the matrix $A = [\lambda_1]$ has the WSP and establishes the base of the induction for the graph G with one vertex. Now assume that our claim holds for $n - 1$, that is, given a graph G_{n-1} on $n - 1$ vertices and $n - 1$ distinct real numbers $\lambda_1, \lambda_2, \dots, \lambda_{n-1}$, there exists a symmetric matrix A_{n-1} whose graph is G_{n-1} , its spectrum is $\{\lambda_1, \lambda_2, \dots, \lambda_{n-1}\}$, and A_{n-1} has the WSP. To prove the claim for n , assume that the vertices of G are labeled by $1, 2, \dots, n$ and let G_{n-1} be the graph obtained from G by removing its n -th vertex. The induction hypothesis applied to G_{n-1} and $\lambda_1, \lambda_2, \dots, \lambda_{n-1}$ yields a matrix A_{n-1} with the desired properties. Now, let

$$(3.2) \quad A = A_{n-1} \oplus [\lambda_n] = \left[\begin{array}{c|c} A_{n-1} & \mathbf{0} \\ \hline \mathbf{0} & \lambda_n \end{array} \right].$$

We will use the Implicit Function Theorem to make the desired entries in the last row and last column of A nonzero, without changing the eigenvalues of the matrix.

Assume that the labeling of the vertices of G is done in such a way that the n -th vertex is adjacent to vertices $1, 2, \dots, d$. Note that A has the WSP by Corollary 2.7, and therefore Lemma 3.1 implies that $J = \text{Jac}_x(g) \Big|_A$ is nonsingular. Thus, by the Implicit Function Theorem, for sufficiently small $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_d$, there exist numbers $\tilde{A}_{11}, \tilde{A}_{22}, \dots, \tilde{A}_{nn}$

close to $A_{11}, A_{22}, \dots, A_{nn}$ such that if

$$\tilde{A} = \left[\begin{array}{cccc|c} \tilde{A}_{11} & A_{12} & \cdots & A_{1,n-1} & \varepsilon_1 \\ A_{21} & \tilde{A}_{22} & \cdots & A_{2,n-1} & \vdots \\ \vdots & \vdots & \ddots & \vdots & \varepsilon_d \\ A_{n-1,1} & A_{n-1,2} & \cdots & \tilde{A}_{n-1,n-1} & 0 \\ \hline \varepsilon_1 & \cdots & \varepsilon_d & 0 & \cdots & 0 & \tilde{A}_{nn} \end{array} \right],$$

then

$$g(\mathbf{x}, \mathbf{y}) \Big|_{\tilde{A}} = \left(\sum_{i=1}^n \lambda_i, \frac{1}{2} \sum_{i=1}^n \lambda_i^2, \dots, \frac{1}{n} \sum_{i=1}^n \lambda_i^n \right).$$

Furthermore, if ε_i 's are chosen to be nonzero, then G is the graph of \tilde{A} . Thus $f(\mathbf{x}, \mathbf{y}) \Big|_{\tilde{A}} = (\lambda_1, \lambda_2, \dots, \lambda_n)$, that is, the spectrum of \tilde{A} equals Λ . It is evident that this solution is not unique. Now recall that the operator norm of the matrix A , denoted $\|A\|_{\text{op}}$, is defined by

$$\|A\|_{\text{op}} = \sup_{\|\mathbf{v}\|_2=1} \|A\mathbf{v}\|_2.$$

Choosing ε_i 's small enough, we can have

$$(3.3) \quad \|\tilde{A} - A\|_{\text{op}} < \varepsilon$$

for any given ε . It only remains to prove that \tilde{A} has the WSP whenever ε is sufficiently small. This is immediate because A defined as in (3.2) has the WSP by Corollary 2.7, and therefore so does \tilde{A} , provided that ε in (3.3) is sufficiently small to allow an application of Lemma 2.4. \square

Remark 3.3. Note that the proof of Theorem 3.2 enables us to control the norm of \tilde{A} in each step. This will be important in examination of the infinite case of this SISP in the next section.

4. INFINITE λ -SISP

In this section we will prove an analogue of Theorem 3.2 for countably infinite (but not necessarily locally finite) graphs. This is done by taking the limit, in a suitable sense, of the matrices \tilde{A} that were constructed in the proof of Theorem 3.2. Then we will show that this limit has the desired properties.

We begin with recalling a few definitions from the spectral theory. Let T be a bounded operator from a Hilbert space \mathcal{H} to itself. The

spectrum of T , and the *point spectrum* of T (consisting of eigenvalues of T) are defined by

$$\sigma(T) = \{\lambda \in \mathbb{C} \mid T - \lambda I \text{ is not invertible}\},$$

and

$$\sigma_p(T) = \{\lambda \in \mathbb{C} \mid T - \lambda I \text{ is not one-to-one}\},$$

respectively. Clearly, $\sigma_p(T)$ is contained in $\sigma(T)$. When \mathcal{H} is finite-dimensional $\sigma(T) = \sigma_p(T)$, since a linear operator on \mathcal{H} is one-to-one if and only if it is invertible. However, when \mathcal{H} is infinite-dimensional T may have additional elements in its spectrum, and may have no eigenvalues.

In general, $\sigma_p(T)$ is contained in a subset of the spectrum, called the *approximate point spectrum* and denoted $\sigma_{\text{app}}(T)$, that plays an important role in the spectral theory in infinite-dimensional spaces. The notion of approximate point spectrum is obtained by strengthening the notion of a one-to-one operator to an operator that is bounded below. By definition, $\lambda \in \sigma_{\text{app}}(T)$ if $T - \lambda I$ is not bounded below, meaning that there are unit vectors u_n with $\|(T - \lambda I)u_n\| \rightarrow 0$ as n tends to infinity. A point λ with this property is called an *approximate eigenvalue* of T .

Let T be a bounded operator on a Hilbert space. We say that T is a *Fredholm operator* if

- (a) the kernel of T has finite dimension, and
- (b) the range of T has finite codimension.

By a theorem of Atkinson [6, Theorem 5.17], T is Fredholm if and only if T is invertible modulo compact operators in the sense that there exists a bounded operator S such that $I - ST$ and $I - TS$ are compact operators.

We shall need a corollary of the following result about a “continuity property” of the spectrum in the proof of our main theorem (See Chapter V, Theorem 4.10 of [10]).

Theorem 4.1. [10, p. 291] *Let S and T be bounded self-adjoint operators on a Hilbert space with the spectra $\sigma(S)$ and $\sigma(T)$, respectively. Then the Hausdorff distance $d_H(\sigma(S), \sigma(T))$ satisfies*

$$d_H(\sigma(S), \sigma(T)) \leq \|S - T\|_{op}.$$

This theorem immediately implies the following corollary.

Corollary 4.2. *Let $\{T_n\}_{n=1}^{\infty}$ be a sequence of bounded self-adjoint operators on a Hilbert space \mathcal{H} , and assume that $T_n \rightarrow T$ in the operator norm. Then for any $\lambda \in \sigma(T)$ and any neighborhood U of λ , there exists an $N \in \mathbb{N}$ such that U intersects $\sigma(T_n)$ nontrivially for all $n > N$.*

Now we are ready to state and prove our main theorem.

Theorem 4.3 (SISP with data (G, Λ)). *Given an infinite graph G on countably many vertices and a compact, infinite set of real numbers Λ , there exists a self-adjoint operator T on the Hilbert space $\ell^2(\mathbb{N})$ such that*

- (i) *the (approximate point) spectrum of T equals Λ , and*
- (ii) *the (real symmetric) standard matrix of T has graph G .*

Proof. Let $\{\lambda_1, \lambda_2, \dots\}$ denote a countable dense subset of Λ . Suppose the vertices of G are labeled by \mathbb{N} and let $G[\{1, 2, \dots, n\}]$ denote the induced subgraph of G on the first n vertices. Then, by (3.3) in the proof of Theorem 3.2, for any $\varepsilon > 0$ and each $n \in \mathbb{N}$ we can find a matrix \tilde{A}_n with the finite graph $G[\{1, 2, \dots, n\}]$, and the spectrum the finite set of distinct real numbers $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$ such that

$$(4.1) \quad \|\tilde{A}_n \oplus [\lambda_{n+1}] - \tilde{A}_{n+1}\|_{\text{op}} < \frac{\varepsilon}{2^n}.$$

For each n define the bounded linear operator T_n on the Hilbert space of square-summable sequences $\ell^2(\mathbb{N})$ (denoted ℓ^2 hereafter for short) such that

$$M_n = \left[\begin{array}{c|ccc} \tilde{A}_n & & & \\ \hline & \lambda_{n+1} & & \\ & & \lambda_{n+2} & \\ & & & \ddots \end{array} \right]$$

is the matrix representation of T_n with respect to the standard Hilbert basis $\mathfrak{B} = \{\mathbf{e}_1, \mathbf{e}_2, \dots\}$ of ℓ^2 , where the entries of M_n that are not shown are zero. It is clear from the definition that the matrices M_n are isospectral. Additionally, by (4.1), for any $i \in \mathbb{N}$

$$\|M_n \mathbf{e}_i - M_{n+1} \mathbf{e}_i\|_2 < \frac{\varepsilon}{2^n}.$$

Thus, the sequence of partial sums $\{\sum_{k=1}^{n-1} (M_{k+1} \mathbf{e}_i - M_k \mathbf{e}_i)\}_{n=1}^\infty$ is absolutely convergent, and therefore the sequence $\{M_n \mathbf{e}_i\}_{n=1}^\infty$ satisfying

$$M_n \mathbf{e}_i = M_1 \mathbf{e}_i + \sum_{k=1}^{n-1} (M_{k+1} \mathbf{e}_i - M_k \mathbf{e}_i)$$

is convergent in ℓ^2 . Let M denote the matrix whose columns are obtained by this limiting process, that is, M is the matrix that $M \mathbf{e}_i = \lim_{n \rightarrow \infty} M_n \mathbf{e}_i$ for each $i \in \mathbb{N}$. Note that for each $n = 1, 2, \dots$ the graph of \tilde{A}_n is the induced subgraph of G on the first n vertices. Thus,

by construction, G is the graph of M . Our next objective is showing that M is indeed the standard matrix of a bounded linear operator $T: \ell^2 \rightarrow \ell^2$. Observe that

$$\begin{aligned}
\|T_n - T_{n+1}\|_{\text{op}} &= \sup_{\|\mathbf{v}\|_2=1} \|T_n \mathbf{v} - T_{n+1} \mathbf{v}\|_2 \\
&= \sup_{\|\mathbf{v}\|_2=1} \left\| \begin{bmatrix} \tilde{A}_n \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} \\ \lambda_{n+1} v_{n+1} \\ 0 \\ \vdots \\ 0 \end{bmatrix} - \tilde{A}_{n+1} \begin{bmatrix} v_1 \\ \vdots \\ v_{n+1} \end{bmatrix} \right\|_2 \\
&= \sup_{\|\mathbf{v}\|_2=1} \left\| \left(\left[\begin{array}{c|c} \tilde{A}_n & \\ \hline & \lambda_{n+1} \end{array} \right] - \tilde{A}_{n+1} \right) \begin{bmatrix} v_1 \\ \vdots \\ v_{n+1} \end{bmatrix} \right\|_2 \\
&< \frac{\varepsilon}{2^n},
\end{aligned}$$

where in the last line we have used the submultiplicative property of the operator norm and (4.1). This inequality immediately implies that the sequence of partial sums $\{\sum_{k=1}^{n-1} (T_{k+1} - T_k)\}_{n=1}^\infty$ is absolutely convergent in the Banach space of bounded operators $\mathcal{B}(\ell^2)$, and hence the sequence $\{T_n\}_{n=1}^\infty$ satisfying

$$T_n = T_1 + \sum_{k=1}^{n-1} (T_{k+1} - T_k)$$

is convergent to a limit T with respect to the operator norm. Since for each $i \in \mathbb{N}$ we have $T \mathbf{e}_i = \lim_{n \rightarrow \infty} T_n \mathbf{e}_i$ and $T_n \mathbf{e}_i = M_n \mathbf{e}_i$, we conclude that $T \mathbf{e}_i = M \mathbf{e}_i$ and thus M is the standard matrix of T .

It remains to prove that $\sigma(T) = \Lambda$. First, we claim that each $\lambda_i \in \{\lambda_1, \lambda_2, \dots\} \subset \Lambda$ is in the spectrum of T , that is, $T - \lambda_i I$ is not invertible. To see this note that as $n \rightarrow \infty$

$$\|(T - \lambda_i I) - (T_n - \lambda_i I)\|_{\text{op}} = \|T - T_n\|_{\text{op}} \rightarrow 0$$

which shows the existence of noninvertible operators in any neighborhood of $T - \lambda_i I$. Since invertibility is an open property in any unital Banach algebra and in particular in $\mathcal{B}(\ell^2)$, we have $\{\lambda_1, \lambda_2, \dots\} \subset \sigma(T)$, and the claim is proved. By taking closures, this inclusion implies $\Lambda \subset \sigma(T)$, because $\{\lambda_1, \lambda_2, \dots\}$ is dense in Λ and $\sigma(T)$ is closed in \mathbb{R} .

Next, since $T_n \rightarrow T$ in the operator norm and $\sigma(T_n) = \Lambda$ for all n , by Corollary 4.2 we conclude that for any $\lambda \in \sigma(T)$, every neighborhood of λ intersects Λ . Hence the reverse inclusion $\sigma(T) \subset \Lambda$ is also established.

Finally, to complete the proof of item (i) in the statement of the theorem note that the spectrum of any self-adjoint operator equals its approximate point spectrum, and T is clearly self-adjoint. \square

The isolated points of the spectrum of any self-adjoint operator are its eigenvalues, for instance, by an application of Gelfand's continuous functional calculus [6]. Thus the isolated points of Λ are contained in the point spectrum of any solution T —obtained as in the proof of Theorem 4.3 or otherwise—for the SISP with data (G, Λ) . On the other hand, by focusing only on the solutions T obtained as in the proof of Theorem 4.3 we will prove that the set of limit points of the spectrum of T equals the *essential spectrum* of T . This will allow us to show that our solutions of every SISP with the same Λ are *approximately unitarily equivalent*. We will then use this equivalence relation to show that the multiplicity of isolated eigenvalues of the constructed solutions is exactly one.

Definition 4.4. Let T be a bounded self-adjoint operator on a Hilbert space. Then the *essential spectrum* of T , $\sigma_{\text{ess}}(T)$, is defined by

$$(4.2) \quad \sigma_{\text{ess}}(T) = \{\lambda \in \mathbb{C} \mid T - \lambda I \text{ is not a Fredholm operator}\}.$$

The essential spectrum of a self-adjoint operator T can be equivalently defined as the complement of its discrete spectrum. That is,

$$(4.3) \quad \sigma_{\text{ess}}(T) = \sigma(T) \setminus \sigma_{\text{discr}}(T),$$

where the discrete spectrum, denoted $\sigma_{\text{discr}}(T)$, is the set of isolated eigenvalues of T with finite multiplicity.

There is a famous classification result due to Weyl, von Neumann, and Berg (Theorem 4.6 below) involving the essential spectrum of self-adjoint operators and the multiplicity of their isolated eigenvalues. To state it we need the following definition.

Definition 4.5. Two bounded operators T_1 and T_2 on Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 are said to be *approximately unitarily equivalent* (written $T_1 \sim_a T_2$) if there is a sequence of unitary isomorphisms of Hilbert spaces $U_n: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that $T_1 = \lim_{n \rightarrow \infty} U_n^* T_2 U_n$.

It is well-known (see for instance [4]) that if $T_1 \sim_a T_2$, then a sequence of unitary operators $\{U_n\}_{n=1}^{\infty}$ can be chosen such that in addition to $T_1 = \lim_{n \rightarrow \infty} U_n^* T_2 U_n$, one also has that $T_1 - U_n^* T_2 U_n$ belongs to the ideal $\mathcal{K}(\mathcal{H}_1)$ of compact operators.

Theorem 4.6 (Weyl–von Neumann–Berg, see [4, Theorem II.4.4]). *Two self-adjoint operators M and N on separable Hilbert spaces are approximately unitarily equivalent if and only if*

- (i) $\sigma_{\text{ess}}(M) = \sigma_{\text{ess}}(N)$, and
- (ii) $\dim \ker(M - \lambda I) = \dim \ker(N - \lambda I)$ for all $\lambda \in \mathbb{C} \setminus \sigma_{\text{ess}}(M)$.

Now we are ready to prove the following corollaries of Theorem 4.3 concerning the limit points of the spectrum and the approximate unitary equivalence of any two constructed solutions of a given SISP.

Corollary 4.7. *Let T be a bounded self-adjoint operator obtained according to the proof of Theorem 4.3 as a solution to the SISP with data (G, Λ) . Then $\sigma_{\text{ess}}(T) = \Lambda'$, where Λ' is the set of limit points of Λ .*

Proof. Let $T = \lim_{n \rightarrow \infty} T_n$ where $\{T_n\}_{n=1}^{\infty}$ is the sequence of operators defined as in the proof of Theorem 4.3 relative to some countable dense subset $\{\lambda_1, \lambda_2, \dots\}$ of Λ . The containment $\Lambda' \subset \sigma_{\text{ess}}(T)$ is clear from (4.3) and the fact that $\Lambda = \sigma(T)$, as shown in Theorem 4.3. To prove the reverse inclusion, let $\lambda \in \sigma_{\text{ess}}(T)$ be arbitrary. By definition (4.2) and Atkinson’s Theorem [6, Theorem 5.17] $T - \lambda I$ is not invertible “modulo compact operators.” More precisely, if we denote the ideal of compact operators in $\mathcal{B}(\ell^2)$ by $\mathcal{K}(\ell^2)$, then $(T - \lambda I) + \mathcal{K}(\ell^2)$ is not an invertible element of the *Calkin algebra* $\mathcal{B}(\ell^2)/\mathcal{K}(\ell^2)$. Moreover, the equality

$$T - \lambda I = [T - \text{diag}(\lambda_1, \lambda_2, \dots)] + [\text{diag}(\lambda_1, \lambda_2, \dots) - \lambda I]$$

implies that

$$(T - \lambda I) + \mathcal{K}(\ell^2) = \text{diag}(\lambda_1 - \lambda, \lambda_2 - \lambda, \dots) + \mathcal{K}(\ell^2),$$

because $T - \text{diag}(\lambda_1, \lambda_2, \dots)$ is the limit of the finite-rank operators $T_n - \text{diag}(\lambda_1, \lambda_2, \dots)$, and hence it is compact. To finish the proof we only need to observe that $\text{diag}(\lambda_1 - \lambda, \lambda_2 - \lambda, \dots) + \mathcal{K}(\ell^2)$ is non-invertible in the Calkin algebra (if and) only if 0 is a limit point of $\{\lambda_1 - \lambda, \lambda_2 - \lambda, \dots\}$. This is equivalent to saying that λ is a limit point of $\{\lambda_1, \lambda_2, \dots\}$ in which case $\lambda \in \Lambda'$. \square

We mention in passing that if Λ' is a singleton, $\Lambda' = \{\lambda\}$, then $T - \lambda I$ is compact. To see this, let T_n and T be as in the proof of Theorem 4.3. If λ is the only limit point of Λ , then all $T_n - \lambda I$ and consequently their limit in operator norm $T - \lambda I$ are compact operators.

Corollary 4.8. *Let S and T be any two bounded self-adjoint operators obtained according to the proof of Theorem 4.3 as solutions to the SISP*

with data (G_1, Λ) and (G_2, Λ) , respectively. Then S and T are approximately unitarily equivalent. In particular, $T \sim_a \text{diag}(\lambda_1, \lambda_2, \dots)$ for any countable dense subset $\{\lambda_1, \lambda_2, \dots\}$ of Λ .

Proof. Let S_n and T_n be defined as in the proof of Theorem 4.3 such that $S = \lim_{n \rightarrow \infty} S_n$ and $T = \lim_{n \rightarrow \infty} T_n$. Then for all n ,

- $\sigma_{\text{ess}}(S_n) = \sigma_{\text{ess}}(T_n) = \Lambda'$, and
- $\dim \ker(S_n - \lambda I) = \dim \ker(T_n - \lambda I) = 1$ for all $\lambda \in \Lambda \setminus \Lambda'$,

because, by construction, each of S_n and T_n can be realized as the direct sum of an $n \times n$ matrix and an infinite diagonal matrix for which verifying these two conditions is straightforward. Thus Theorem 4.6 implies $S_n \sim_a T_n$. For each n , let $\{U_{n,k}\}_{k=1}^{\infty}$ denote the sequence of unitary operators that satisfy $S_n = \lim_{k \rightarrow \infty} U_{n,k}^* T_n U_{n,k}$ and in addition, by passing to a subsequence of $\{U_{n,k}\}_{k=1}^{\infty}$ if necessary, we can arrange that when $k = 1$ we have

$$\|S_n - U_{n,1}^* T_n U_{n,1}\|_{\text{op}} < \max \left\{ \|T_n - T\|_{\text{op}}, \|S_n - S\|_{\text{op}}, \frac{1}{n} \right\}.$$

Define the sequence of unitary operators $\{V_n\}_{n=1}^{\infty}$ by setting $V_n = U_{n,1}$. We wish to prove that $S = \lim_{n \rightarrow \infty} V_n T V_n^*$. This is accomplished by a 3ε -argument as follows. Given $\varepsilon > 0$, choose $N > 0$ large enough such that

$$\max \left\{ \|T_n - T\|_{\text{op}}, \|S_n - S\|_{\text{op}}, \frac{1}{n} \right\} < \varepsilon$$

whenever $n > N$. Then for any such n ,

$$\begin{aligned} \|S - V_n^* T V_n\|_{\text{op}} &\leq \|S - S_n\|_{\text{op}} + \|S_n - V_n^* T_n V_n\|_{\text{op}} + \|V_n^* T_n V_n - V_n^* T V_n\|_{\text{op}} \\ &= \|S - S_n\|_{\text{op}} + \|S_n - U_{n,1}^* T_n U_{n,1}\|_{\text{op}} + \|T_n - T\|_{\text{op}} \\ &< 3 \max \left\{ \|T_n - T\|_{\text{op}}, \|S_n - S\|_{\text{op}}, \frac{1}{n} \right\} \\ &< 3\varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, our claim is proved. \square

“Approximate unitary equivalence” cannot be replaced by the stronger notion of “unitary equivalence” in Corollary 4.8 as the following simple example shows. Let G be the empty graph on countably many vertices and let Λ be the compact set $\{\frac{1}{n} \mid n \in \mathbb{N}\} \cup \{0\}$. Then our construction in the proof of Theorem 4.3 produces the following solutions for the SISP with data (G, Λ) :

$$(4.4) \quad \text{diag} \left(1, \frac{1}{2}, \frac{1}{3}, \dots \right) \text{ and } \text{diag} \left(0, 1, \frac{1}{2}, \dots \right)$$

corresponding to two different choices of countable dense subsets of Λ , namely $\Lambda \setminus \{0\}$ and Λ . The two operators in (4.4) are approximately unitarily equivalent but clearly not unitarily equivalent.

Remark 4.9. Corollary 4.8 immediately implies that if T is a solution of the SISP with data (G_1, Λ_1) constructed as above, then any isolated eigenvalue of T has multiplicity one. To see this it suffices to let G_2 be the empty graph on countably many vertices and let $\Lambda_2 = \Lambda_1$ so that a solution to the SISP with data (G_2, Λ_2) is the diagonal operator $S = \text{diag}(\lambda_1, \lambda_2, \dots)$ for a countable dense subset $\{\lambda_1, \lambda_2, \dots\}$ of Λ . Since $S \sim_a T$ by Corollary 4.8, an application of Theorem 4.6 to the pair of self-adjoint operators S and T gives

$$\dim \ker(T - \lambda I) = 1 \text{ for all } \lambda \in \sigma(T) \setminus \sigma_{\text{ess}}(T) = \Lambda \setminus \Lambda'.$$

5. CONCLUSION

Motivated by some problems in science and engineering such as the inverse Sturm-Liouville problem, in this manuscript we have introduced a structured inverse spectrum problem (SISP) for infinite graphs and have shown the existence of a (non-unique) solution for this problem. We have used the Jacobian method to establish a procedure for controlling the norm of the solutions of the finite problem in a way that the constructed sequence of solutions for finite cases converges to a solution of the infinite problem. We have also shown that any two solutions of the SISP constructed by our method are approximately unitarily equivalent.

In an extension of this work [11] we relax the compactness condition on Λ in the main result of this paper by working with unbounded operators. More precisely, given an infinite graph G on countably many vertices, and a closed, infinite set Λ of real numbers, we prove the existence of an unbounded self-adjoint operator whose graph is G and whose spectrum is Λ .

6. ACKNOWLEDGEMENT

The research of the first author is supported by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- [1] G. E. Backus and J. F. Gilbert. Numerical applications of a formalism for geophysical inverse problems. *Geophysical Journal of the Royal Astronomical Society*, 13:247–276, 1967.

- [2] Wayne W. Barrett, Shaun M. Fallat, H. Tracy Hall, Leslie Hogben, Jephian C.-H. Lin, and Bryan L. Shader. Generalizations of the Strong Arnold Property and the minimum number of distinct eigenvalues of a graph. *Electron. J. Comb.*, 24:P2.40 (28 pages), 2017.
- [3] Göran Borg. Eine umkehrung der Sturm-Liouvilleschen eigenwertaufgabe. *Acta Math.*, 78:1–96, 1946.
- [4] Kenneth R. Davidson. *C*-Algebras by Example*. American Mathematical Society, Providence, R.I., 1996.
- [5] Carl de Boor and Gene H. Golub. The numerically stable reconstruction of a Jacobi matrix from spectral data. *Linear Algebra and Its Applications*, 21:245–260, 1978.
- [6] Ronald G. Douglas. *Banach Algebra Techniques in Operator Theory*. Graduate Texts in Mathematics 179. Springer-Verlag, New York, second edition, 1998.
- [7] António Leal Duarte. Construction of acyclic matrices from spectral data. *Linear Algebra and its Applications*, 113:173–182, 1989.
- [8] Ole H. Hald. The inverse Sturm-Liouville problem and the Rayleigh-Ritz method. *Mathematics of Computation*, 32:687–705, 1978.
- [9] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 2nd edition, 2013.
- [10] Tosio Kato. *Perturbation theory for linear operators*. Springer-Verlag, New York, 1995.
- [11] Ehssan Khanmohammadi. A structured inverse spectrum problem for infinite graphs and unbounded operators. *arXiv:1708.01708 [math.SP]*.
- [12] Mark G. Krein. Determination of the density of a non-homogeneous symmetric cord by its frequency spectrum. *Dokl. Akad. Nauk SSSR*, 76:345–348, 1951.
- [13] Flavia Lenti, Ferdinando Nunziata, Claudio Estatico, and Maurizio Migliaccio. Analysis of reconstructions obtained solving ℓ^p -penalized minimization problems. *IEEE Transactions on Geoscience and Remote Sensing*, 1:1–11, 2015.
- [14] Ian G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1995.
- [15] Keivan Hassani Monfared and Bryan L. Shader. Construction of matrices with a given graph and prescribed interlaced spectral data. *Linear Algebra and its Applications*, 438:4348–4358, 2013.
- [16] William F. Stokey. Vibration of systems having distributed mass and elasticity. In Allan Piersol and Thomas Paez, editors, *Harris' shock and vibration handbook, 6th edition*, chapter 7, pages 7.1–7.50. McGraw-Hill Education, 2009.

UNIVERSITY OF CALGARY
E-mail address: k1monfared@gmail.com

FRANKLIN & MARSHALL COLLEGE
E-mail address: ehssan@fandm.edu