

ECE-2025

Spring-05

LECTURE #2

Phase & Time-Shift

Complex Exponentials

14-Jan-05

2/25/2005

ECE-2025 Spring-2005 JMc

1

INFORMATION

- **MATLAB: 6pm Tues, Wed in BH-216**
- LABS start NEXT week (**Tuesday**)
 - Attend correct section (in Bunger-Henry 216)
 - ECE Computer acct password:
 - last 4 digits of gtID# + "!" + 3 character month of birth + "\$" + 4 digit year of birth + "."
 - For example, if your gtID# was 012345678 and you were born 1/1/1970
5678!Jan\$1970.
(<http://www.ece-help.gatech.edu/labs/overview.html#account>)
- RECITATIONS
 - **Bring Calculator to Recitation next week**
 - **Complex arithmetic**
 - Attend your assigned time
 - **ACTIVE** participation

2/25/2005

ECE-2025 Spring-2005 JMc

2

Web-CT Info

- Check the Bulletin Board for msgs
 - **MAKE YOUR OWN POSTINGS**
- Web-CT Login: everyone should have acct
- PDF Files on WebCT
 - Lectures are being posted (4 per page)
 - Lab #1 will be posted soon
 - Get PDF file of Lab #1 from WebCT
 - Hard copy of Instructor Verification Sheet
 - HW #1 was posted as PDF
 - HW #1 due next week
 - HW #2 will be posted soon

2/25/2005

ECE-2025 Spring-2005 JMc

3

Homework Info

- HWs will be posted on Friday/Sat
 - Covered in Rec during the following Week
 - Due the week after that (9+ days later)
- **Format** info on WebCT
 - Cover page for Homework
- Prob Set #1 due **in RECITATION**
 - **At the beginning of class**
 - Solutions will be posted to WebCT after the last Recitation on Thursday which ends at 6pm

2/25/2005

ECE-2025 Spring-2005 JMc

4


Homework Formatting

- Include a Cover page with
 - Name
 - Lab section, ie, L05, L20, etc.
 - Recitation Prof's name
 - **Download example from Web-CT**
- Write on **ONE** side only
 - Use Engineer's paper or plain white paper
- STAPLE

2/25/2005

ECE-2025 Spring-2005 JMc

5



2/25/2005

JUMP

6

READING ASSIGNMENTS

- This Lecture:
 - Chapter 2, Sects. 2-3 to 2-5
- Appendix A: Complex Numbers
- Appendix B: MATLAB
- Next Lecture: finish Chap. 2,
 - Section 2-6 to end

2/25/2005

ECE-2025 Spring-2005 JMc

7

LECTURE OBJECTIVES

- Define Sinusoid Formula from a plot
- Relate TIME-SHIFT to PHASE

Introduce an **ABSTRACTION**:
Complex Numbers **represent** Sinusoids
Complex Exponential Signal

$$z(t) = X e^{j\omega t}$$

2/25/2005

ECE-2025 Spring-2005 JMc

8

SINUSOIDAL SIGNAL

$$A \cos(\omega t + \varphi)$$

- **FREQUENCY** ω
 - Radians/sec
 - or, Hertz (cycles/sec)
 - $\omega = (2\pi) f$
- **PERIOD** (in sec)
 $T = \frac{1}{f} = \frac{2\pi}{\omega}$
- **AMPLITUDE** A
 - Magnitude
- **PHASE** φ

2/25/2005

ECE-2025 Spring-2005 JMc

9

PLOTTING COSINE SIGNAL from the FORMULA

$$5 \cos(0.3\pi t + 1.2\pi)$$

- Determine **period**:
 $T = 2\pi / \omega = 2\pi / 0.3\pi = 20/3$
- Determine a **peak** location by solving
 $(\omega t + \varphi) = 0$
- **Peak at t=-4**

2/25/2005


ECE-2025 Spring-2005 JMc

10

ANSWER for the PLOT

$$5 \cos(0.3\pi t + 1.2\pi)$$

- Use $T=20/3$ and the peak location at $t=-4$



TIME-SHIFT

- In a mathematical formula we can replace t with $t-t_m$
$$x(t - t_m) = A \cos(\omega(t - t_m))$$
- Then the $t=0$ point moves to $t=t_m$
- Peak value of $\cos(\omega(t-t_m))$ is now at $t=t_m$


2/25/2005

ECE-2025 Spring-2005 JMc

12

TIME-SHIFTED SINUSOID

$$x(t+4) = 5 \cos(0.3\pi(t+4)) = 5 \cos(0.3\pi(t - (-4)))$$



2/25/2005

ECE-2025 Spring-2005 JMc

13

PHASE \leftrightarrow TIME-SHIFT

- Equating the formulas:

$$A \cos(\omega(t - t_m)) = A \cos(\omega t + \phi)$$

- and we obtain:

$$-\omega t_m = \phi$$

- or,

$$t_m = -\frac{\phi}{\omega}$$

2/25/2005

ECE-2025 Spring-2005 JMc

14

SINUSOID from a PLOT

- Measure the period, T
 - Between peaks or zero crossings
- Compute frequency: $\omega = 2\pi/T$
- Measure time of a peak: t_m
 - Compute phase: $\phi = -\omega t_m$
- Measure height of positive peak: A


3 steps

2/25/2005

ECE-2025 Spring-2005 JMc

15

(A, ω, ϕ) from a PLOT



$$T = \frac{0.01 \text{ sec}}{1 \text{ period}} = \frac{1}{100}$$

$$\omega = \frac{2\pi}{T} = \frac{2\pi}{0.01} = 200\pi$$

$$t_m = -0.00125 \text{ sec}$$


$$\phi = -\omega t_m = -(200\pi)(-0.00125) = 0.25\pi$$

2/25/2005

ECE-2025 Spring-2005 JMc

16

SINE DRILL (MATLAB GUI)



2/25/2005

17

PHASE is AMBIGUOUS

- The cosine signal is periodic

- Period is 2π

$$A \cos(\omega t + \varphi + 2\pi) = A \cos(\omega t + \varphi)$$

- Thus adding any multiple of 2π leaves $x(t)$ unchanged

if $t_m = \frac{-\varphi}{\omega}$, then

$$t_{m_2} = \frac{-(\varphi + 2\pi)}{\omega} = \frac{-\varphi}{\omega} - \frac{2\pi}{\omega} = t_m - T$$


2/25/2005

ECE-2025 Spring-2005 JMc

18

COMPLEX NUMBERS

- To solve: $z^2 = -1$
 - $z = j$
 - Math and Physics use $z = i$
- Complex number: $z = x + jy$




Cartesian
coordinate
system

2/25/2005

ECE-2025 Spring-2005 JMc

19


PLOT COMPLEX NUMBERS



2/25/2005

20

COMPLEX ADDITION = VECTOR Addition




*** POLAR FORM ***

Vector Form

- Length = 1
- Angle = θ

Common Values

- j has angle of 0.5π
- -1 has angle of π
- $-j$ has angle of 1.5π
- also, angle of $-j$ could be $-0.5\pi = 1.5\pi - 2\pi$
- because the PHASE is **AMBIGUOUS**




POLAR <--> RECTANGULAR

- Relate (x,y) to (r,theta)

$$r^2 = x^2 + y^2$$

$$\theta = \text{Tan}^{-1}\left(\frac{y}{x}\right)$$



$$x = r \cos \theta$$

$$y = r \sin \theta$$


Most calculators do Polar-Rectangular

Need a notation for POLAR FORM

Euler's FORMULA

Complex Exponential

- Real part is cosine
- Imaginary part is sine
- Magnitude is one



$$e^{j\theta} = \cos(\theta) + j \sin(\theta)$$


$$re^{j\theta} = r \cos(\theta) + jr \sin(\theta)$$

COMPLEX EXPONENTIAL

$$e^{j\omega t} = \cos(\omega t) + j \sin(\omega t)$$

Interpret this as a **Rotating Vector**

- $\theta = \omega t$
- Angle changes vs. time
- ex: $\omega = 20\pi$ rad/s
- Rotates 0.2π in 0.01 secs



$$e^{j\theta} = \cos(\theta) + j \sin(\theta)$$

2/25/2005

ECE-2025 Spring-2005 JMc

25

COS = REAL PART

Real Part of Euler's

$$\cos(\omega t) = \Re\{e^{j\omega t}\}$$

General Sinusoid

$$x(t) = A \cos(\omega t + \varphi)$$

So,

$$\begin{aligned} A \cos(\omega t + \varphi) &= \Re\{A e^{j(\omega t + \varphi)}\} \\ &= \Re\{A e^{j\varphi} e^{j\omega t}\} \end{aligned}$$

2/25/2005

ECE-2025 Spring-2005 JMc

26

REAL PART EXAMPLE

$$A \cos(\omega t + \varphi) = \Re\{A e^{j\varphi} e^{j\omega t}\}$$

Evaluate:

$$x(t) = \Re\{-3j e^{j\omega t}\}$$

Answer:

$$\begin{aligned} x(t) &= \Re\{(-3j) e^{j\omega t}\} \\ &= \Re\{3e^{-j0.5\pi} e^{j\omega t}\} = 3 \cos(\omega t - 0.5\pi) \end{aligned}$$

2/25/2005

ECE-2025 Spring-2005 JMc

27

COMPLEX AMPLITUDE

General Sinusoid

$$x(t) = A \cos(\omega t + \varphi) = \Re\{A e^{j\varphi} e^{j\omega t}\}$$

Complex AMPLITUDE = X

$$z(t) = X e^{j\omega t} \quad X = A e^{j\varphi}$$

Then, any Sinusoid = REAL PART of $X e^{j\omega t}$

$$x(t) = \Re\{X e^{j\omega t}\} = \Re\{A e^{j\varphi} e^{j\omega t}\}$$

2/25/2005

ECE-2025 Spring-2005 JMc

28