

Math 2551 Final Review Fall 2022

1. Find \mathbf{T} , \mathbf{N} , and the curvature κ for $\mathbf{r}(t) = 3\sin(t)\mathbf{i} + 3\cos(t)\mathbf{j} + 4t\mathbf{k}$.
2. Write acceleration in terms of its tangential and normal components for $\mathbf{r}(t) = \langle t + 1, 2t, t^2 \rangle$ at $t = 1$.
3. Find the limit: $\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2 - y^2 + 5}{x^2 + y^2 + 2}$
4. Find all second-order partial derivatives for $w = x \sin(x^2y)$
5. Evaluate $\frac{dw}{dt}$ for $w = 2ye^x - \ln(z)$ if $x(t) = \ln(t^2 + 1)$, $y = \arctan(t)$, $z = e^t$ at $t = 1$.
6. Find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ for the surface $z^3 - xy + yz + y^3 - 2 = 0$ at the point $(1, 1, 1)$.
7. Find the derivative of the function $f(x, y) = 2xy - 3y^2$ in the direction $4\mathbf{i} + 3\mathbf{j}$ at the point $(5, 5)$.
8. Find the tangent plane and normal line at $(2, 0, 2)$ of the surface $2z - x^2 = 0$.
9. Find all local maxima, local minima, and saddle points for $f(x, y) = x^2 + xy + y^2 + 3x - 3y + 4$.
10. Find the absolute maximum and minimum of $f(x, y) = 2x^2 - 4x + y^2 - 4y + 1$ on the closed triangular region bounded by the lines $x = 0$, $y = 2$, and $y = 2x$.
11. Find the maximum value of $f(x, y) = 49 - x^2 - y^2$ on the line $x + 3y = 10$.
12. Write an iterated integral using (a) vertical cross-sections and (b) horizontal cross-sections whose region of integration is the region bounded by $y = \sqrt{x}$, $y = 0$, $x = 9$.
13. Sketch the region of integration and evaluate the integral $\int_0^1 \int_0^{y^2} 3y^3 e^{xy} dx dy$.
14. Change to polar coordinates and evaluate the integral $\int_{-1}^1 \int_0^{\sqrt{1-x^2}} dy dx$.
15. Integrate the function $f(x, y) = 3 - 4x$ over the region below $z = 4 - xy$ and above the rectangle $0 \leq x \leq 2$, $0 \leq y \leq 1$ in the xy -plane.
16. Find the volume of the region that lies inside the sphere $x^2 + y^2 + z^2 = 2$ and outside the cylinder $x^2 + y^2 = 1$.
17. Use the change of coordinates $x(u, v) = \frac{u}{v}$, $y(u, v) = uv$ to evaluate the integral $\iint_R \left(\sqrt{\frac{y}{x}} + \sqrt{xy} \right) dA$, where R is the region in the first quadrant bounded by $xy = 1$, $xy = 0$, $y = x$, and $y = 4x$.
18. Evaluate the line integral $\int_C (xy + y + z) ds$ along the curve $\mathbf{r}(t) = 2t\mathbf{i} + t\mathbf{j} + (2 - 2)\mathbf{k}$ for $0 \leq t \leq 1$.
19. Find the flow of the field $\mathbf{F} = \langle -4xy, 8y, 2 \rangle$ along the curve $\mathbf{r}(t) = \langle t, t^2, 1 \rangle$, $0 \leq t \leq 2$.
20. Find the counterclockwise circulation and the outward flux of the field $\mathbf{F} = x\mathbf{i} + y\mathbf{j}$ around/through the unit circle centered at the origin.
21. Find the potential function for $\mathbf{F} = e^{y+2z}(\mathbf{i} + x\mathbf{j} + 2x\mathbf{k})$.
22. Use Green's Theorem to find counterclockwise circulation and outward flux of the field $\mathbf{F} = \langle y^2 - x^2, x^2 + y^2 \rangle$ for the curve C enclosing the region bounded by $y = 0$, $x = 3$, and $y = x$.
23. Use a parameterization to write a double integral for the area of the surface S which is the portion of the cone $z = 2\sqrt{x^2 + y^2}$ between the planes $z = 2$ and $z = 6$.

24. Evaluate $\iint_S 2y \, d\sigma$ over the surface S which is the part of the cylinder $y^2 + z^2 = 4$ between $x = 0$ and $x = 3 - z$.
25. Let S be the surface that consists of the part of the paraboloid $z = 4 - x^2 - y^2$ above the xy -plane and below the cone $z = 3\sqrt{x^2 + y^2}$.
- Sketch S .
 - Find a parameterization of S .
 - Calculate the area of S .
26. Let S be the surface consisting of the top half $z \geq 0$ of the sphere $x^2 + y^2 + z^2 = 9$, together with the disk $x^2 + y^2 \leq 9, z = 0$, its base in the xy -plane. Use the divergence theorem to evaluate

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma,$$

where $\mathbf{F}(x, y, z) = 3xy^2\mathbf{i} + 3x^2y\mathbf{j} + z^3\mathbf{k}$.

27. Let S be the part of the surface $z = 4x^2 + y^2 - 4$ beneath the plane $z = 5$. Let C be the bounding curve of S in the plane $z = 5$, traversed counterclockwise and suppose S is oriented accordingly (normals towards the z -axis). Let $\mathbf{F}(x, y, z) = \langle 2y, 4x, e^x \rangle$. Use Stokes' Theorem to evaluate the curl integral

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\sigma.$$

28. Let S be the surface of the cylinder defined by $y^2 + z^2 = 4$ between the planes $x = -1$ and $x = 3$. Let $\mathbf{F}(x, y, z) = e^{xy}\mathbf{i} + e^x y\mathbf{j} + e^x z\mathbf{k}$.
- Sketch S .
 - Find a parameterization of S .
 - Let \mathbf{n} be an outward pointing unit normal for S . Evaluate

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma$$

by direct calculation (do not use the Divergence Theorem).