




Delaware Trends in Adolescent Sexual Risk Behaviors

February 8, 2011

From the 2009 Youth Risk Behavior Survey
University of Delaware Center for Drug and
Alcohol Studies




Percentage of students who had ever been physically forced to have sexual intercourse when they did not want to



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Percentage of students who ever had sexual intercourse



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Percentage of students who had sexual intercourse for the first time before age 13 years



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Percentage of students who had sexual intercourse with four or more people during their life



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Percentage of students who had sexual intercourse with one or more people during the past three months



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Among students who had sexual intercourse during the past three months, the percentage who drank alcohol or used drugs before last sexual intercourse



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Among students who had sexual intercourse during the past three months, the percentage who used a condom during the last sexual intercourse



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware




Among students who had sexual intercourse during the past three months, the percentage who used birth control pills to prevent pregnancy before last sexual intercourse



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware

Percentage of students who had ever been taught in school about AIDS or HIV infection



Note: * : Denotes linear significant change
 † : Denotes quadratic significant change

Source: 2009 Delaware Youth Risk Behavior Survey, Center for Drug and Alcohol Studies, University of Delaware