



Note

The minimum degree distance of graphs of given order and size

Orest Bucicovschi, Sebastian M. Cioabă*

Department of Mathematics, University of California, San Diego, La Jolla, CA 92093-0112, United States

ARTICLE INFO

Article history:

Received 24 January 2008

Received in revised form 13 March 2008

Accepted 28 March 2008

Available online 8 May 2008

Keywords:

Degree

Distance

Wiener index

ABSTRACT

In this note, we study the degree distance of a graph which is a degree analogue of the Wiener index. Given n and e , we determine the minimum degree distance of a connected graph of order n and size e .

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

Our graph notation is standard (see [11]). Let $\mathcal{G}_{n,e}$ denote the family of connected graphs on n vertices and e edges and $\mathcal{G}_n = \cup_e \mathcal{G}_{n,e}$. For $G \in \mathcal{G}_{n,e}$, let $d(x, y)$ denote the distance between vertices x and y and let $d(x)$ denote the degree of x . Define $D(x) = \sum_{y \in V(G)} d(x, y)$ and

$$\begin{aligned} D'(G) &= \sum_{x \in V(G)} d(x)D(x) = \sum_{x \in V(G)} d(x) \sum_{y \in V(G)} d(x, y) \\ &= \frac{1}{2} \sum_{x, y \in V(G)} d(x, y)(d(x) + d(y)). \end{aligned}$$

The parameter $D'(G)$ is called the degree distance of G and it was introduced by Dobrynin and Kochetova [4] and Gutman [6] as a weighted version of the Wiener index $D(G) = \sum_{x, y \in V(G)} d(x, y)$. Actually, when G is a tree on n vertices, these parameters are closely related as $D'(G) = 2D(G) - n(n - 1)$ in this case (see [6]).

In [9], Ioan Tomescu confirmed a conjecture of Dobrynin and Kochetova from [4] and proved the following result.

Theorem 1.1. For $n \geq 2$,

$$\min_{G \in \mathcal{G}_n} D'(G) = 3n^2 - 7n + 4. \quad (1)$$

Also, $K_{1, n-1}$ is the only connected graph G on n vertices such that $D'(G) = 3n^2 - 7n + 4$.

Recently, Alexandru Ioan Tomescu [10] determined the minimum value of $D'(G)$ when $G \in \mathcal{G}_{n,e}$ and $e \in \{n, n + 1\}$. In this note, we give a short proof of Theorem 1.1 and we extend the results of [10] for all n and e such that $1 \leq n - 1 \leq e$.

For k a real number, let $\Sigma_k(G)$ denote the sum of the k -th powers of the degrees of G . This parameter has been studied in various contexts in [2,3,5,7]. We denote by $\sigma_2(n, e)$ the maximum value of $\Sigma_2(G)$ when G is a graph (not necessarily connected) with n vertices and e edges. Also, let $\Sigma_2(n, e) = \max_{G \in \mathcal{G}_{n,e}} \Sigma_2(G)$. We discuss the connection between these two parameters in Section 3. This connection is already apparent in the next theorem which is our main result.

* Corresponding author. Tel.: +1 858 534 2623; fax: +1 858 534 5273.

E-mail addresses: obucicov@math.ucsd.edu (O. Bucicovschi), scioaba@math.ucsd.edu (S.M. Cioabă).

Theorem 1.2. *If $1 \leq n - 1 \leq e$ and G is a connected graph with n vertices and e edges, then*

$$D'(G) \geq 4(n - 1)e - \Sigma_2(n, e) = 4(n - 2)e - n^2 + 5n - 4 - \sigma_2(n - 1, e - n + 1).$$

Equality happens if and only if G is a join of K_1 and a graph G' on $n - 1$ vertices and $e - n + 1$ edges with $\Sigma_2(G') = \sigma_2(n - 1, e - n + 1)$.

In Section 2, we give a short proof of Theorem 1.1. In Section 3, we present the proof of Theorem 1.2.

2. A short proof of Theorem 1.1

In this section we give a short proof of equality (1).

Proof. For $x \in V(G)$, we have that

$$\begin{aligned} D(x) &= \sum_{y \in V(G)} d(x, y) = \sum_{y: d(x, y)=1} d(x, y) + \sum_{y: d(x, y) \geq 2} d(x, y) \\ &\geq 1 \cdot d(x) + 2 \cdot (n - d(x) - 1) = 2n - 2 - d(x). \end{aligned}$$

It follows that

$$\begin{aligned} D'(G) &= \sum_{x \in V(G)} d(x)D(x) \geq \sum_{x \in V(G)} d(x)(2n - 2 - d(x)) \\ &= \sum_{x \in V(G)} d(x)(2n - 3) - \sum_{x \in V(G)} d(x)(d(x) - 1) \\ &= 2e(2n - 3) - \sum_{x \in V(G)} d(x)(d(x) - 1) \\ &\geq 2e(2n - 3) - \sum_{x \in V(G)} (n - 1)(d(x) - 1) \\ &= 2e(2n - 3) - (n - 1)(2e - n) = 2e(n - 2) + (n - 1)n. \end{aligned}$$

Since $e \geq n - 1$, this proves inequality (1). Equality happens if and only if G has $n - 1$ edges (which means G is a tree) and $d(x) = n - 1$ for each vertex x with $d(x) > 1$. This is equivalent to G being isomorphic to $K_{1, n-1}$ which finishes the proof. \square

3. Proof of Theorem 1.2

In this section, we show how to find the minimum $D'(G)$ over all connected graphs G with n vertices and e edges.

Proof. From the proof of the previous theorem, we obtain that for any vertex $x \in V(G)$,

$$D'(x) \geq (2n - 2)d(x) - d^2(x).$$

Summing up over all $x \in V(G)$, we get

$$\begin{aligned} D'(G) &= \sum_{x \in V(G)} D'(x) \geq \sum_{x \in V(G)} ((2n - 2)d(x) - d^2(x)) \\ &= 4(n - 1)e - \Sigma_2(G). \end{aligned}$$

Since $\Sigma_2(G) \leq \Sigma_2(n, e)$, the first inequality follows immediately. Equality happens if and only the diameter of G is 2 and $\Sigma_2(G) = \Sigma_2(n, e)$.

We claim that if G is a connected graph on n vertices and e edges such that $\Sigma_2(G) = \Sigma_2(n, e)$, then G contains a vertex of degree $n - 1$.

Assume otherwise and let u be a vertex of G of maximum degree with $d(u) < n - 1$. This implies that there exist vertices v and w such that $uv, vw \in E(G)$, but $uw \notin E(G)$. Consider the graph H obtained from G by adding the edge uw and removing the edge vw . It is easy to see that $H \in \mathcal{G}_{n, e}$. Also, we have that

$$\begin{aligned} \Sigma_2(H) - \Sigma_2(G) &= (d(u) + 1)^2 + (d(v) - 1)^2 - d^2(u) - d^2(v) \\ &= 2(d(u) - d(v)) + 2 \geq 2. \end{aligned}$$

This inequality contradicts the fact that $\Sigma_2(G) = \Sigma_2(n, e)$ and proves our claim.

Thus, the equality condition is equivalent to G being a join of K_1 and a graph G' on $n - 1$ vertices and $e - n + 1$ edges and $\Sigma_2(G) = \Sigma_2(n, e)$. Note that the graph G' is not necessarily connected.

A simple calculation yields that

$$\begin{aligned}\Sigma_2(G) &= (n-1)^2 + \sum_{x \in V(G')} d^2(x) \\ &= (n-1)^2 + \sum_{x \in V(G')} ((d(x)-1)^2 + 2(d(x)-1) + 1) \\ &= (n-1)^2 + \sum_{x \in V(G')} (d(x)-1)^2 + 2 \sum_{x \in V(G')} d(x) - (n-1) \\ &= (n-1)^2 + \Sigma_2(G') + 2(2e - n + 1) - (n-1) \\ &= n^2 - 5n + 4 + 4e + \Sigma_2(G').\end{aligned}$$

This implies that $\Sigma_2(G') = \sigma_2(n-1, e-n+1)$ as well as that

$$\Sigma_2(n, e) = n^2 - 5n + 4 + 4e + \sigma_2(n-1, e-n+1). \quad (2)$$

The proof of Theorem 1.1 is now complete. \square

When $e = n$ or $e = n + 1$, we obtain the main results of [10].

Given n and $e \geq n - 1$, we have seen that determining $\Sigma_2(n, e)$ is equivalent to finding $\sigma_2(n - 1, e - n + 1)$. Ahlswede and Katona [1] have determined $\sigma_2(n - 1, e - n + 1)$ and the extremal graphs attaining this bound. This problem has a long history and its solution is nontrivial (see [1,8] for more details). For the sake of completeness, we outline the result of Ahlswede and Katona below.

For a given order p and size $q \leq \binom{p}{2}$, the *quasi-complete* graph C_p^q is the graph with vertex set $\{1, \dots, p\}$ and q edges defined as follows: i is adjacent to j for any $i, j \in \{1, \dots, a\}$ and $a + 1$ is adjacent to $1, \dots, b$ where a and b are the unique integers such that $q = \binom{a}{2} + b$ and $0 \leq b < a$. The *quasi star* S_p^q is the complement of the $C_p^{\binom{p}{2}-q}$.

In [1], Ahlswede and Katona showed that

$$\begin{aligned}\sigma_2(p, q) &= \max(\Sigma_2(C_p^q), \Sigma_2(S_p^q)) \\ &= \max(2q(a-1) + b(b+1), (n(n-1) - 2q)(a-1) + b(b+1) + 4q(n-1) - (n-1)^2n).\end{aligned}$$

for any p and q with $q \leq \binom{p}{2}$. They also proved that

$$\sigma_2(p, q) = \begin{cases} \Sigma_2(S_p^q), & \text{if } 0 \leq q \leq \frac{\binom{p}{2}}{2} - p \\ \Sigma_2(C_p^q), & \text{if } \frac{\binom{p}{2}}{2} + p \leq q \leq \binom{p}{2} \end{cases}$$

Ahlswede and Katona also showed that in the range $\frac{\binom{p}{2}}{2} - p < q < \frac{\binom{p}{2}}{2} + p$, it is more difficult to determine which one of C_p^q or S_p^q attains the maximum. More precisely, they proved that there are infinitely many p 's for which $\sigma_2(p, q) = \Sigma_2(S_p^q)$ for all $q < \frac{1}{2} \binom{p}{2}$ and $\sigma_2(p, q) = \Sigma_2(C_p^q)$ for all $q > \frac{1}{2} \binom{p}{2}$. Also, there are infinitely many p 's for which the previous statement is not true (see [1], Lemma 8).

In [9], Ioan Tomescu disproved a conjecture from [4] and showed that

$$\frac{n^4}{27} + O(n^3) \leq \max_{G \in \mathcal{G}_n} D'(G) \leq \frac{2n^4}{27} + O(n^3). \quad (3)$$

Tomescu conjectured that the lower bound is the exact value of $\max_{G \in \mathcal{G}_n} D'(G)$. This conjecture seems difficult at present time.

Acknowledgments

The authors dedicate this note to Professor Ioan Tomescu on the occasion of his 65-th birthday. The authors thank the referee for her or his comments. The second author's research is partially supported by an NSERC postdoctoral fellowship.

References

- [1] R. Ahlswede, G.O.H. Katona, Graphs with maximal number of adjacent pairs of edges, Acta Math. Acad. Sci. Hungar. 32 (1978) 97–120.
- [2] B. Bollobás, V. Nikiforov, Degree powers in graphs with forbidden subgraphs, Electron. J. Combin. 11 (2004) R42.
- [3] S.M. Cioabă, Sums of powers of the degrees of a graph, Discrete Math. 306 (2006) 1959–1965.
- [4] A.A. Dobrynin, A.A. Kochetova, Degree distance of a graph: A degree analogue of the Wiener index, J. Chem. Inf. Comput. Sci. 34 (1994) 1082–1086.
- [5] Z. Füredi, A. Kündgen, Moments of graphs in monotone families, J. Graph Theory 51 (2006) 37–48.
- [6] I. Gutman, Selected properties of the Schulz molecular topological index, J. Chem. Inf. Comput. Sci. 34 (1994) 1087–1089.

- [7] Y. Hu, X. Li, Y. Shi, T. Xu, Connected (n, m) -graphs with minimum and maximum zeroth-order general Randić index, *Discrete Appl. Math.* 155 (2007) 1044–1054.
- [8] V. Nikiforov, The sum of squares of the degrees: Sharp asymptotics, *Discrete Math.* 307 (2007) 3187–3193.
- [9] I. Tomescu, Some extremal properties of the degree distance of a graph, *Discrete Appl. Math.* 98 (1999) 159–163.
- [10] A.I. Tomescu, Unicyclic and bicyclic graphs having minimum degree distance, *Discrete Appl. Math.* 156 (2008) 125–130.
- [11] D.B. West, *Introduction to Graph Theory*, second ed., Prentice Hall, 2001.