

Series: Long-lasting Impact of Early-Life Stress and Adversity

## Opinion

# Neurodevelopmental Optimization after Early-Life Adversity: Cross-Species Studies to Elucidate Sensitive Periods and Brain Mechanisms to Inform Early Intervention

Joan L. Luby,<sup>1,\*</sup> Tallie Z. Baram,<sup>2</sup> Cynthia E. Rogers,<sup>1</sup> and Deanna M. Barch<sup>1,3,4</sup>

**Human brain development is influenced by early-life experiences, particularly during sensitive periods, with impact on cognitive and emotional outcomes. Understanding how the timing and nature of such experiences (including adversity, trauma, and enrichment) govern their influence on brain organization is crucial for harnessing key environmental factors early in life to enhance brain development. Here we synthesize findings from human and animal studies focusing on sensitive periods and their regional and circuit specificity and highlight the challenge and power of such cross-species approaches in informing the ‘next steps’ to optimize cognitive and emotional health in developing children. We propose designs for neurodevelopmental optimization research programs utilizing randomized enhancement trials in early childhood to inform public health strategies on prevention and early intervention.**

## Early Experiences and Vulnerability to, or Protection against, Mental Illness

The finding that human brain development is strongly influenced by the environment, particularly during early-life sensitive periods, has important global mental and public health implications. An increasing body of empirical data from human and animal studies demonstrates strong associations (in humans) and causal relations (in animals) between early environmental factors and brain maturation [1–7]. Early-life experiences are known to play an important role in influencing cognitive and emotional outcomes in humans through their impact on neurodevelopment. There is also robust evidence for the strong effects of early adversity on risk for psychopathology [8]. Conversely, a large body of work demonstrates the central importance of early-life nurturance for healthy social and emotional development [9,10]. Building on this work, we address how empirically informed timing of preventive or developmentally enhancing interventions may be used to achieve larger and more sustained neuroprotective effects against the negative consequences of adversity. To accomplish this, a more comprehensive understanding of how environmental factors influence specific aspects of the complex machinery of brain development is required. Data informing these processes from animal, human, and cross-species studies would facilitate the harnessing of modifiable factors early in life to support healthier brain development as part of a proactive practice of optimizing child development [11,12].

We propose here a program of research as the foundation for a new science of neurodevelopmental optimization. Building on causal inferences and mechanistic data from the animal-model literature, we posit that there is a need to obtain a nuanced empirical picture of how the timing and type of adverse exposures impacts specific aspects of neurodevelopment and the correlated cognitive and emotional capabilities in young children. We highlight the

## Highlights

Human brain development is influenced by exposure to early-life experiences, including enrichment and adversity, with cognitive and emotional consequences including vulnerability to or protection from mental illness.

The timing of the exposure is critical, because there are sensitive periods when vulnerability is augmented; sensitive periods may pertain also to the timing of enrichment or mitigation efforts.

Animal studies indicate that sensitive periods are specific for distinct brain regions and circuits, providing a timing framework for selective and insult-specific interventions.

Capitalizing on findings gleaned from animal-model studies and human imaging in childhood, we propose the ‘next steps’ in cross-species research towards the goal of optimizing cognitive and emotional health in developing children.

Neurodevelopmental optimization research should address issues including deprivation, unpredictability, and insecure attachment as well as, potentially sleep, diet, and the gut microbiome via carefully timed randomized enhancement trials.

<sup>1</sup>Department of Psychiatry, Washington University School of Medicine, Early Emotional Development Program, 4444 Forest Park Avenue, St. Louis, MO, USA  
<sup>2</sup>Departments of Pediatrics, Anatomy/Neurobiology, and Neurology, University of California, Irvine, Irvine, CA 92697, USA



potential of caregiving interventions that enhance warmth, sensitivity, and predictability as well as child cognitive and emotional skill-building programs delivered at key sensitive periods. These strategies carry a promise to nourish neurodevelopment, mitigate the risk for psychopathology, and enhance human potential.

<sup>3</sup>Department of Psychological and Brain Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA

<sup>4</sup>Department of Radiology, Washington University School of Medicine, St. Louis, MO, USA

### Key Considerations for Neurodevelopmental Optimization: Timing and Specificity

Information about the nature, timing, regional specificity, and mechanisms of environmental factors that influence the trajectory of brain maturation is necessary to enable neurodevelopmental optimization programs. Timing relates to whether there are sensitive periods in the development of specific brain regions and circuits subserving specific emotional and cognitive functions, during which children might be either particularly vulnerable to environmental adversity or receptive to enhancement. Regional specificity refers to whether environmental effects are broad, related to general factors that influence brain development as a whole, or more specific to particular brain regions and circuits. Identifying regional specificity will inform the nature of enhancements targeting specific regions or circuits and their cognate functions. The specificity of environmental factors refers to the degree to which different types of early adversity (e.g., trauma/abuse, deprivation, unpredictability/chaos, poverty) and/or enhancement (caregiver support, cognitive stimulation, enriched diet or sleep) have similar or different effects on brain development both globally and regionally. Empirical mapping of these patterns is necessary to inform the type and timing of the most effective optimization strategies.

\*Correspondence:  
lubyj@wustl.edu (J.L. Luby).

### Sensitive Periods across Species

Whereas the brain is influenced by the environment through experience-dependent processes throughout the lifespan, sensitive periods are windows of time during which the brain is especially susceptible to these influences [13–15]. Sensitive periods allow the brain architecture to be maximally informed by experience to optimize function for events expected later in life and are well documented in both the basic neuroscience and the child development literature [13–16]. Notably, the timing and duration of sensitive periods are themselves experience dependent [14–17].

Importantly, it is becoming evident that different brain regions and circuits have distinct trajectories of development and sensitive periods [16–19]. This information is critical to enable the translation of ground-breaking experimental studies in animals to human interventions. Whereas older work compares phases of total brain growth across species, newer studies avoid assigning a global brain age to rodents that is then equated with human age. Rather, the maturation of specific brain circuits and regions is compared [18,19]. For example, for hippocampal formation development, the developmental state of a human full-term neonate might be equated with that of a 5–7 day Sprague Dawley rat, with infancy encompassing the second week of life in the rodent [18]. A similar approach to identifying homologs in brain age across humans and rodents has been recently employed for the reward circuit, including the ventral striatum/nucleus accumbens, the ventral tegmental area, and interconnected amygdala nuclei and cortical regions, again suggesting homology between the middle first postnatal week in the rodent and the human neonate (details see Table 1 in [19]).

Sensitive periods for specific regions and circuits in animals and to a lesser extent in humans are being delineated. Such studies attest to the importance of the timing of early-life experiences, because the ages of sensitive periods for distinct regions or circuits differ [15–17,20]. For example, the sensitive period for the effects of light signals on visual system organization in kittens spans the first postnatal weeks [20]. In humans, an ‘amblyopic’ or otherwise deprived eye during the first postnatal months provokes enduring loss of normal function in the primary visual cortex

and life-long deficits in vision (amblyopia), suggesting homologous timing-sensitive plasticity processes during sensitive periods across species [14].

The sensitive period for the patterns of tones on the tonotopic organization of the auditory cortex in the mouse involves postnatal days 7–14. In rodents, the sensitive period for early-life adversity seems to include the first two postnatal weeks, influencing both the maturation of specific brain circuits and functional outcomes [6,21–29]. Deprivation from maternal signals [30,31] or chaotic unpredictable patterns of care [23–28] promote vulnerability to memory and emotional deficits.

Whereas it is not possible to directly translate sensitive periods across species, it is helpful to consider that at least some of the neurobiological mechanisms that generate sensitive periods are likely to be common across mammals [14–16,20]: the organization of brain circuits involves the generation of synaptic connections, followed by their strengthening and persistence or their pruning and elimination. It is generally believed that the first step, the recognition by pre- and post-synaptic elements of their future ‘partners’, is genetically encoded and relatively insensitive to the environment. The second stage, involving activity-dependent processes and molecular triggers such as the maturation of specific neurotransmitter systems, influences persistent versus eliminated synapses and constitutes the sensitive period. Because the timing of this second phase can be estimated in humans and rodents from the developmental trajectory of each circuit in each species, the relative timing of sensitive periods can be estimated across species [32,33].

Artificial augmentation of maternal care via ‘handling’ has been widely shown to enhance cognitive and emotional outcomes in rodents. Daily brief separations predictably promote recurrent intense barrages of maternal care behavior on the return of the pups to the cage [34], and this enrichment has consistently been shown to lead to a well-regulated response to stress as well as enhanced memory functions [35–37], as found in other enrichment studies in animals [34,38–41]. Accordingly, natural variation in the quantity and quality of maternal care behaviors correlates with pups’ outcomes, supporting the positive effects of extensive and consistent maternally derived sensory input to developing rat pups on cognitive and social behaviors [22,42].

In humans, the landmark Bucharest Early Intervention Project (BEIP) randomized institutionalized children to therapeutic foster homes and compared them with those remaining in institutional care [15]. Results suggested that the first 2 years of life might be the developmental period most sensitive to the negative effects of primary caregiver deprivation on later cognitive and emotional outcomes, a finding augmented by a second sample that suggests that different brain circuits have different sensitive periods, emphasizing the amygdala–prefrontal cortex circuitry [11]. However, much more work is needed to address sensitive periods in human development, and we propose the use of randomized controlled trials of discrete and targeted enhancements in early childhood that are informed by animal studies as the next most feasible and important scientific step.

### Specificity of Experience Type for Neurodevelopment

Evidence of some neural specificity of types of adversity in both animals and humans is available. These types of adversity include abuse, neglect, deprivation, poverty, and unpredictability and fragmentation of parental care and environmental signals. The broader construct of adverse childhood experiences (ACEs) includes many of these factors as well as exposure to parental mental disorder and criminal behavior [43]. Notably, these forms of early adversity often co-occur and share enhanced risk for poor neurodevelopmental outcomes and psychiatric disorders [18,23,44–47]. A critical issue is whether forms of early adversity converge on the same aspects of brain structure and function or whether there is evidence of neural specificity to particular forms

of adversity. The challenge in dissociating these diverse components of adversity in human studies led to assessment of the issue in experimental animals, where paradigms have been designed to simulate distinct aspects of early-life adversity. These include separation from the dam once or chronically [30,31,48–50] (for a review see [24]) and simulated poverty/resource scarcity [6,23,25–29]. Notably, as is the case in human adversity, most of these paradigms intermingle, generating maternal stress that disrupts maternal care patterns rendering them unpredictable or abusive [6,21,23,26] (for reviews see [24,29]). Thus, assessing the selective contributions of different components of adversity to cognitive and emotional outcomes remains a significant challenge in both human and animal studies.

### Specificity: Which Brain Regions Are Impacted and When?


The functional consequences of early-life adversity are a result of disruption of the development of the underlying brain regions and circuits. To date, much of the literature has focused on particular brain regions, but future work will need to more clearly embed such regions in the larger networks in which they function. Many studies converge on a relation between distinct types of early adversity and hippocampal structure and function, including reductions in hippocampal volume associated with poverty [51–59], reduced maternal support [60,61], and abuse/ACE [1,4,62]. There is also evidence for reduced amygdala volume associated with poverty [52,53,55,56,63], which may vary by age [64]. Alterations in striatum structure in relation to early adversity, often associated with deficits in reward processing, have also been reported [65,66]. Controlled animal work supports the causal nature of such associations [11,21,25,28,30,49,67].

A longitudinal neuroimaging study in humans found more complex developmentally specific interactions between the timing of experience (preschool, school age, adolescence) and both positive and negative regional brain effects [44]. Specifically, interactions between preschool ACEs and school age maternal support were found for both hippocampus and amygdala volumes, such that school-age maternal support was associated with greater volumes only in the context of low preschool ACEs. However, for the caudate, a pattern suggesting early emerging additive reductions in caudate volume was associated independently with preschool maternal support and ACEs that were stable over time. These findings suggest that there is regional, and likely circuit, specificity to the timing of adversity and support as they influence brain maturation, providing clues for the design of future neurodevelopmental optimization strategies.

### Neurodevelopmental Enhancement Programs Informed by Timing and Specificity

We aim to employ this empirical knowledge of distinct sensitive periods and generate additional information to achieve larger and more sustained neuroprotective effects against the negative consequences of adversity (Figure 1). To achieve this, we propose the use of focused enrichment paradigms in randomized controlled trials in early childhood. These studies should design enhancement interventions building on known sensitive periods in animal models and emerging human work and apply them to young child samples. The use of environmental enhancement that targets parenting more broadly is an important and feasible strategy that could test the importance of protections or enhancements during sensitive periods at varying ages and most importantly targeting birth to age 5 years. In addition, the application of enhancements that directly target the child and augment specific emotional or cognitive skill building at different ages will elucidate sensitive periods for human cognitive and emotional development. Specifically, infants/young children facing a variety of forms of adversity can be randomized to usual care versus enriched or stimulating settings for periods of time or enhanced parenting at certain age periods, ideally those shown to be sensitive to particular inputs (e.g., before age 2 years compared with later in preschool for enhanced parenting). Another design would be to expose young children

to intensive training for specific cognitive and emotion skills (e.g., emotion recognition, executive function) also at specific developmental periods and then compare them with those who do not receive the training. We suggest targeting interventions for children living in poverty or facing adversity who have primary caregivers whose support can be harnessed for such interventions as a first step. We propose to employ empirically validated early mental health interventions. One such example with large effect sizes and enduring efficacy and high feasibility is Attachment and Bio-behavioral Catch Up (ABC) [68] that is focused on enhancing early attachment. Other effective programs include child–parent psychotherapy (CPP) [69], video-based intervention to promote positive parenting (MIPP) [70], and several forms of the preschool intervention Parent–Child Interaction Therapy (PCIT) that have also demonstrated large effect sizes and enduring efficacy [71,72]. These interventions can be tested at different age periods and varying ‘doses’ with the



Legend

-  Adversity
-  Protection/nurturance/predictability

**Figure 1. Factors in the Optimization of Early Childhood Neurodevelopment.** The schematic illustrates the theoretical optimization model, where umbrellas represent the specific need for protection from adversity at key timepoints (to be empirically determined) based on sensitive periods. Enhancement is applied during phases of development also based on these empirically determined periods and the child’s individual needs. Lightning bolts represent adversity that developing children may face. The placement of these icons in the figure is currently speculative, awaiting empirical anchoring based on animal and human enhancement trials to inform the timing of sensitive periods.

effects on brain structure and function pre- and postintervention assessed. To assess effects on brain network organization, one could employ, for instance, resting-state neuroimaging during sleep, and this could be augmented (or replaced) by feasible and less costly measures of neural function such as task-based electroencephalography (EEG) or event-related potentials. The proposed studies will inform and provide the building blocks of a new neurodevelopmental optimization approach that is pragmatic and cost-effective and could be applied broadly in public health settings.

Other broad and overall underexplored targets for the enhancement of child neurodevelopment include sleep, diet, and the gut microbiome. Early-life adversity may disrupt these targets, which may contribute to suboptimal brain development. For example, the role of the developmental timing and quality of sleep and circadian rhythms during neurodevelopment should be further studied, as well as the possible effects of their disruption. Diet acting on brain development either directly or via alterations of the gut microbiome has increasingly been a focus of research, and emerging evidence in animal models suggests that replenishing specific micro- and macro-nutrients early in life may mitigate the cognitive consequences of experimentally imposed early-life adversity [73–75]. The effects of all of these modifiable environmental factors on neurodevelopment should be further clarified in terms of the nature and timing of exposures, as they represent potential pathways for timing- and context-dependent neurodevelopmental optimization.

### Concluding Remarks

We propose a concept- and data-driven approach to neurodevelopmental optimization to enable enhancement programs promoting optimal cognitive and emotional outcomes in early childhood (Figure 1). Given the greater focus to date on risk factors, relatively little attention has been given to the notion of optimization, which can be useful to those at both high and low risk. For these programs to be effective, they need to incorporate basic principles of brain development and build on experimental animal models that enable establishment of causality and mechanisms (see Outstanding Questions). The proposed interventional studies in humans could provide the foundation for large-scale, cost-effective preventive approaches to mental illness.

### Acknowledgments

Supported by National Institute of Mental Health (NIMH) grants RO1 MH R01MH090786, P50 MH06889, RO1 MH73136, and R01MH113570.

### References

- Bick, J. and Nelson, C.A. (2016) Early adverse experiences and the developing brain. *Neuropsychopharmacology* 41, 177
- Klengel, T. and Binder, E.B. (2015) Epigenetics of stress-related psychiatric disorders and gene-environment interactions. *Neuron* 86, 1343–1357
- Farah, M.J. (2018) Socioeconomic status and the brain: prospects for neuroscience-informed policy. *Nat. Rev. Neurosci.* 19, 428–438
- McLaughlin, K.A. et al. (2019) Childhood adversity and neural development: a systematic review. *Annu. Rev. Dev. Psychol.* 1, 277–312
- Short, A.K. and Baram, T.Z. (2019) Early-life adversity and neurological disease: age-old questions and novel answers. *Nat. Rev. Neurol.* 15, 657–669
- Chen, Y. and Baram, T.Z. (2016) Toward understanding how early-life stress reprograms cognitive and emotional brain networks. *Neuropsychopharmacology* 41, 197
- Wang, X.-D. and Schmidt, M.V. (2016) Molecular mechanisms for reprogramming hippocampal development and function by early-life stress. *Front. Mol. Neurosci.* 9, 6
- McLaughlin, K.A. et al. (2012) Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *Arch. Gen. Psychiatry* 69, 1151–1160
- Egeland, B. et al. (1993) Resilience as process. *Dev. Psychopathol.* 5, 517–528
- Luthar, S.S. and Eisenberg, N. (2017) Resilient adaptation among at-risk children: harnessing science toward maximizing salutary environments. *Child Dev.* 88, 337–349
- Callaghan, B.L. and Tottenham, N. (2016) The neuro-environmental loop of plasticity: a cross-species analysis of parental effects on emotion circuitry development following typical and adverse caregiving. *Neuropsychopharmacology* 41, 163–176
- Davis, E.P. et al. (2017) Exposure to unpredictable maternal sensory signals influences cognitive development across species. *Proc. Natl. Acad. Sci. U. S. A.* 114, 10390–10395
- Gabard-Dumam, L.J. and McLaughlin, K.A. (2019) Do sensitive periods exist for exposure to adversity? *Biol. Psychiatry* 85, 789–791
- Knudsen, E.J. (2004) Sensitive periods in the development of the brain and behavior. *J. Cogn. Neurosci.* 16, 1412–1425
- Reh, R.K. et al. (2020) Critical period regulation across multiple timescales. *Proc. Natl. Acad. Sci. U. S. A.* Published online June 5, 2020. <https://doi.org/10.1073/pnas.1820836117>

### Outstanding Questions

When – during development – are the key sensitive periods for social, emotional, and cognitive skills?

Are there specific sensitive periods and regional specificities for the effects of different types of adversity and nurturance on neurodevelopment?

To apply neurobiological information from experimental animal models to humans, focused and concerted transdisciplinary cross-species studies are needed. Can research programs be designed to enhance productive and cutting-edge cross-species research?

How can findings from both experimental models and human research be harnessed to design a neurodevelopmental enhancement program to protect children from adversities at key time periods and inform the timing of developmental enhancement in specific domains?

Can insights from such research efforts be applied on a broad, public health level for prevention and early intervention strategies?

16. Hensch, T.K. and Bilimoria, P.M. (2012) Re-opening windows: manipulating critical periods for brain development. *Cerebrum* 2012, 11
17. Huh, C.Y. *et al.* (2020) Long-term monocular deprivation during juvenile critical period disrupts binocular integration in mouse visual thalamus. *J. Neurosci.* 40, 585–604
18. Avishai-Eliner, S. *et al.* (2002) Stressed-out, or in (utero)? *Trends Neurosci.* 25, 518–524
19. Birnie, M.T. *et al.* (2020) Plasticity of the reward circuitry after early life adversity: mechanisms and significance. *Biol. Psychiatry* 87, 875–884
20. Espinosa, J.S. and Stryker, M.P. (2012) Development and plasticity of the primary visual cortex. *Neuron* 75, 230–249
21. Bolton, J.L. *et al.* (2018) Anhedonia following early-life adversity involves aberrant interaction of reward and anxiety circuits and is reversed by partial silencing of amygdala corticotropin-releasing hormone gene. *Biol. Psychiatry* 83, 137–147
22. Champagne, F.A. (2010) Early adversity and developmental outcomes: Interaction between genetics, epigenetics, and social experiences across the life span. *Perspect. Psychol. Sci.* 5, 564–574
23. Molet, J. *et al.* (2016) Fragmentation and high entropy of neonatal experience predict adolescent emotional outcome. *Transl. Psychiatry* 6, e702
24. van Bodegom, M. *et al.* (2017) Modulation of the hypothalamic-pituitary-adrenal axis by early life stress exposure. *Front. Cell. Neurosci.* 11, 87
25. Molet, J. *et al.* (2016) MRI uncovers disrupted hippocampal microstructure that underlies memory impairments after early-life adversity. *Hippocampus* 26, 1618–1632
26. Raineki, C. *et al.* (2019) During infant maltreatment, stress targets hippocampus, but stress with mother present targets amygdala and social behavior. *Proc. Natl. Acad. Sci. U. S. A.* 116, 22821–22832
27. Bath, K.G. *et al.* (2017) Early life stress leads to developmental and sex selective effects on performance in a novel object placement task. *Neurobiol. Stress* 7, 57–67
28. Lucassen, P.J. *et al.* (2013) Perinatal programming of adult hippocampal structure and function; emerging roles of stress, nutrition and epigenetics. *Trends Neurosci.* 36, 621–631
29. Walker, C.-D. *et al.* (2017) Chronic early life stress induced by limited bedding and nesting (LBN) material in rodents: critical considerations of methodology, outcomes and translational potential. *Stress* 20, 421–448
30. Huot, R.L. *et al.* (2002) Neonatal maternal separation reduces hippocampal mossy fiber density in adult Long Evans rats. *Brain Res.* 950, 52–63
31. Kosten, T.A. *et al.* (2007) Memory impairments and hippocampal modifications in adult rats with neonatal isolation stress experience. *Neurobiol. Learn. Mem.* 88, 167–176
32. Schafer, D.P. *et al.* (2012) Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron* 74, 691–705
33. Zeanah, C.H. *et al.* (2011) VI. Sensitive periods. *Monogr. Soc. Res. Child Dev.* 76, 147–162
34. Korosi, A. and Baram, T.Z. (2009) The pathways from mother's love to baby's future. *Front. Behav. Neurosci.* 3, 27
35. Plotsky, P.M. and Meaney, M.J. (1993) Early, postnatal experience alters hypothalamic corticotropin-releasing factor (CRF) mRNA, median eminence CRF content and stress-induced release in adult rats. *Mol. Brain Res.* 18, 195–200
36. Singh-Taylor, A. *et al.* (2018) NRSF-dependent epigenetic mechanisms contribute to programming of stress-sensitive neurons by neonatal experience, promoting resilience. *Mol. Psychiatry* 23, 648–657
37. Korosi, A. *et al.* (2010) Early-life experience reduces excitation to stress-responsive hypothalamic neurons and reprograms the expression of corticotropin-releasing hormone. *J. Neurosci.* 30, 703–713
38. Francis, D.D. *et al.* (2002) Environmental enrichment reverses the effects of maternal separation on stress reactivity. *J. Neurosci.* 22, 7840–7843
39. Bredy, T.W. *et al.* (2004) Peripubertal environmental enrichment reverses the effects of maternal care on hippocampal development and glutamate receptor subunit expression. *Eur. J. Neurosci.* 20, 1355–1362
40. Sotnikov, S. *et al.* (2014) Enriched environment impacts trimethylthiazoline-induced anxiety-related behavior and immediate early gene expression: critical role of C rhr1. *Eur. J. Neurosci.* 40, 2691–2700
41. Curley, J.P. *et al.* (2009) Social enrichment during postnatal development induces transgenerational effects on emotional and reproductive behavior in mice. *Front. Behav. Neurosci.* 3, 25
42. Parent, C.I. and Meaney, M.J. (2008) The influence of natural variations in maternal care on play fighting in the rat. *Dev. Psychobiol.* 50, 767–776
43. Harris, N.B. (2018) *The Deepest Well: Healing the Long-Term Effects of Childhood Adversity*, Houghton Mifflin Harcourt
44. Luby, J.L. *et al.* (2019) Association of timing of adverse childhood experiences and caregiver support with regionally specific brain development in adolescents. *JAMA Netw. Open* 2, e1911426
45. Gee, D.G. *et al.* (2013) Early developmental emergence of human amygdala-prefrontal connectivity after maternal deprivation. *Proc. Natl. Acad. Sci. U. S. A.* 110, 15638–15643
46. Goff, B. *et al.* (2013) Reduced nucleus accumbens reactivity and adolescent depression following early-life stress. *Neuroscience* 249, 129–138
47. Davis, E.P. *et al.* (2019) Across continents and demographics, unpredictable maternal signals are associated with children's cognitive function. *EBioMedicine* 46, 256–263
48. González-Pardo, H. *et al.* (2019) Influence of environmental enrichment on the volume of brain regions sensitive to early life stress by maternal separation in rats. *Psicothema* 31, 46–52
49. Guijarro, J.Z. *et al.* (2007) Effects of brief and long maternal separations on the HPA axis activity and the performance of rats on context and tone fear conditioning. *Behav. Brain Res.* 184, 101–108
50. Spencer-Booth, Y. and Hinde, R.A. (1971) The effects of 13 days maternal separation on infant rhesus monkeys compared with those of shorter and repeated separations. *Anim. Behav.* 19, 595–605
51. Luby, J.L. *et al.* (2013) The effects of poverty on childhood brain development: the mediating effect of caregiving and stressful life events. *JAMA Pediatr.* 167, 1135–1142
52. Hanson, J.L. *et al.* (2015) Behavioral problems after early life stress: contributions of the hippocampus and amygdala. *Biol. Psychiatry* 77, 314–323
53. Jednorog, K. *et al.* (2012) The influence of socioeconomic status on children's brain structure. *PLoS One* 7, e42486
54. Hanson, J.L. *et al.* (2011) Association between income and the hippocampus. *PLoS One* 6, e18712
55. Noble, K.G. *et al.* (2012) Hippocampal volume varies with educational attainment across the life-span. *Front. Hum. Neurosci.* 6, 307
56. Butterworth, P. *et al.* (2012) The association between financial hardship and amygdala and hippocampal volumes: results from the PATH through life project. *Soc. Cogn. Affect. Neurosci.* 7, 548–556
57. Piras, F. *et al.* (2011) Education mediates microstructural changes in bilateral hippocampus. *Hum. Brain Mapp.* 32, 282–289
58. Staff, R.T. *et al.* (2012) Childhood socioeconomic status and adult brain size: childhood socioeconomic status influences adult hippocampal size. *Ann. Neurol.* 71, 653–660
59. Palacios-Barrios, E.E. and Hanson, J.L. (2019) Poverty and self-regulation: connecting psychosocial processes, neurobiology, and the risk for psychopathology. *Compr. Psychiatry* 90, 52–64
60. Luby, J.L. *et al.* (2016) Preschool is a sensitive period for the influence of maternal support on the trajectory of hippocampal development. *Proc. Natl. Acad. Sci. U. S. A.* 113, 5742–5747
61. Luby, J.L. *et al.* (2012) Maternal support in early childhood predicts larger hippocampal volumes at school age. *Proc. Natl. Acad. Sci. U. S. A.* 109, 2854–2859
62. Herzog, J.I. and Schmahl, C. (2018) Adverse childhood experiences and the consequences on neurobiological, psychosocial, and somatic conditions across the lifespan. *Front. Psychiatry* 9, 420

63. Noble, K.G. *et al.* (2012) Neural correlates of socioeconomic status in the developing human brain. *Dev. Sci.* 15, 516–527
64. Merz, E.C. *et al.* (2018) Socioeconomic status, amygdala volume, and internalizing symptoms in children and adolescents. *J. Clin. Child Adolesc. Psychol.* 47, 312–323
65. Novick, A.M. *et al.* (2018) The effects of early life stress on reward processing. *J. Psychiatr. Res.* 101, 80–103
66. Dennison, M.J. *et al.* (2019) Differential associations of distinct forms of childhood adversity with neurobehavioral measures of reward processing: a developmental pathway to depression. *Child Dev.* 90, e96–e113
67. Guadagno, A. *et al.* (2018) Morphological and functional changes in the preweaning basolateral amygdala induced by early chronic stress associate with anxiety and fear behavior in adult male, but not female rats. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 81, 25–37
68. Dozier, M. and Bernard, K. (2019) *Coaching Parents of Vulnerable Infants: The Attachment and Biobehavioral Catch-Up Approach*, Guilford
69. Lieberman, A.F. (2004) Child–parent psychotherapy: a relationship-based approach to the treatment of mental health disorders in infancy and early childhood. In *Treating Parent–Infant Relationship Problems: Strategies for Intervention* (Sameroff, A.J. *et al.*, eds), pp. 97–122, Guilford
70. Van Zeijl, J. *et al.* (2006) Attachment-based intervention for enhancing sensitive discipline in mothers of 1-to 3-year-old children at risk for externalizing behavior problems: a randomized controlled trial. *J. Consult. Clin. Psychol.* 74, 994
71. Fisher, P.A. *et al.* (2016) Promoting healthy child development via a two-generation translational neuroscience framework: the Filming Interactions to Nurture Development video coaching program. *Child Dev. Perspect.* 10, 251–256
72. Eyberg, S.M. *et al.* (2001) Parent–Child Interaction Therapy with behavior problem children: one and two year maintenance of treatment effects in the family. *Child Fam. Behav. Ther.* 23, 1–20
73. Naninck, E.F. *et al.* (2017) Early micronutrient supplementation protects against early stress-induced cognitive impairments. *FASEB J.* 31, 505–518
74. Abbink, M.R. *et al.* (2020) The effects of early life stress, postnatal diet modulation, and long-term Western-style diet on later-life metabolic and cognitive outcomes. *Nutrients* 12, 570
75. Yam, K.-Y. *et al.* (2019) Increasing availability of  $\omega$ -3 fatty acid in the early-life diet prevents the early-life stress-induced cognitive impairments without affecting metabolic alterations. *FASEB J.* 33, 5729–5740