

Cognitive Control and Schizophrenia

Psychological and Neural Mechanisms

Deanna M. Barch and Todd S. Braver

Schizophrenia is a complex and debilitating psychiatric disorder that affects approximately one percent of the population. Lay conceptions of schizophrenia typically focus on symptoms such as hallucinations, delusions, and disorganized speech, which are often considered the hallmark features of this disorder. However, clinicians, researchers, and theorists have long noted that individuals with schizophrenia also commonly suffer from disturbances in memory and cognition, often severely so. Interestingly, recent research suggests that disturbances in social and occupational functioning in individuals with schizophrenia may be more influenced by the severity of their cognitive deficits than the severity of symptoms such as hallucinations and delusions (Green, Kern, Braff, & Mintz, 2000). Such findings have led to a resurgence of interest in identifying the nature of cognitive abnormalities in schizophrenia. A close examination of the types of symptoms and cognitive disturbances displayed by individuals suggests that many of these disturbances appear to reflect an inability to control or regulate their cognitive and emotional states. In this chapter, we review the evidence that one of the core cognitive disturbances in schizophrenia is a deficit in one or more components of executive function, which leads to disturbances in the ability to appropriately regulate thoughts and behavior in accordance with internal goals. More specifically, we suggest that individuals with schizophrenia suffer from a disturbance in a specific type of executive control process that we refer to as a deficit in the ability to represent and maintain context.

Researchers have long recognized that individuals with schizophrenia appear to show profound deficits on cognitive tasks that are thought to require what is collectively referred to as "executive functions" or cognitive control functions. For example, numerous studies have shown that individuals with schizophrenia are impaired on tasks, such as the Wisconsin Card Sorting Task, that require a number of different components of executive function, including problem solving, set switching, and working

memory (Weinberger, Berman, & Zec, 1986). At the same time, individuals with schizophrenia also display deficits in a number of other cognitive domains, including selective attention, inhibition, working memory, perceptual integration (Silverstein, Kovacs, Corry, & Valone, 2000), and sustained attention (Carter, Robertson, & Nordahl, 1992; Chapman & McGhie, 1962; Cornblatt & Keilp, 1994; Gold, Carpenter, Randolph, Goldberg, & Weinberger, 1997; Gold, Randolph, Carpenter, Goldberg, & Weinberger, 1992; Nuechterlein & Dawson, 1984; Park & Holzman, 1992). These findings of deficits in a broad array of task domains raised the question of whether individuals with schizophrenia have a number of independent (or at least semi-independent) cognitive deficits, or whether there is some fundamental or basic component of cognitive control that is impaired in this disorder, which in turn contributes to disturbances on tasks thought to measure many different cognitive domains. In prior work, we and our colleagues have put forth the hypothesis that one of the fundamental disturbances in cognitive control present in schizophrenia is a deficit in the ability to represent and maintain context information (Barch et al., 2001; Braver, Barch, & Cohen, 1999a; Braver & Cohen, 1999; Cohen & Servan-Schreiber, 1992) because of a disturbance in the function of dopamine in dorsolateral prefrontal cortex.

To develop more explicit theories and testable predictions about context processing deficits in schizophrenia, we have drawn upon computational modeling as a tool for specifying the neural mechanisms that support context processing (e.g., dorsolateral prefrontal cortex and the dopamine system) and how specific disturbances to these mechanisms lead to cognitive impairments (Braver, 1997; Braver et al., 1999a; Braver, Barch, & Cohen, 1999b; Braver & Cohen, 1999; Braver, Cohen, & McClelland, 1997; Braver, Cohen, & Servan-Schreiber, 1995; Cohen & Servan-Schreiber, 1992). These models were constructed within the parallel distributed processing (PDP), or "neural network" framework, allowing the quantitative simulation of human performance in cognitive tasks using principles of processing that are similar to those believed to apply in the brain (McClelland, 1993; Rumelhart & McClelland, 1986). The nature of these models and the results of simulations are discussed in detail elsewhere (Braver, 1997; Braver et al., 1999a, 1999b; Braver & Cohen, 1999; Braver et al., 1997; Braver et al., 1995).

CONTEXT PROCESSING

As discussed in many of the other chapters in this volume, such as that by West and Bowry, cognitive control requires the ability to detect, adjust, and respond to changing contingencies and feedback in the environment and no doubt requires a number of different functions and mechanisms. In our own work, we focused on several such mechanisms that we think are important for cognitive control, including context representation, context

intenance, context updating, conflict detection, and subgoal processing (Barch et al., 2001; Barch et al., 2001; Botvinick, Braver, Barch, Carter, & Cohen, 2001; Braver et al., 1999a; Braver & Bongiolatti, 2002; Braver, Cohen, Barch, 2002; O'Reilly, Noelle, Braver, & Cohen, 2002). In regards to understanding cognitive control deficits in schizophrenia, we argued that deficits in the representation, maintenance, and updating of context may be central to this disorder. By context information, we mean information that must be "actively" held in mind in a form that allows it to be used to mediate appropriate behavior. In our models, representations of context are used specifically to support task-relevant information against sources of interference that can occur simply as a function of time (e.g., noise accumulating in the system) or because of specific competing processes (e.g., the need to process other stimuli, distractions). Context representations can comprise a variety of different types of information, such as a specific prior stimulus, the result of processing a sequence of prior stimuli, or more abstract information such as task instructions. As an example, take the following sentence: "In order to keep pigs, you need a pen." In this sentence, the first clause serves as context that biases you toward the appropriate meaning of the word "pen" (a fenced enclosure) for this sentence, rather than the more common meaning of the word "pen" (a writing instrument). This is a case in which the result of processing the first part of the sentence (e.g., a sequence of prior stimuli) creates a contextual representation that can bias future behavior (e.g., semantic interpretation of the word "pen"). As another example, take the well-known Stroop task. In one condition of this task, participants are shown color words written in different colors (e.g., the word "RED" written in blue). In the absence of any other information, a participant's natural response is to read the word rather than to name the color, as this is what one typically does when presented with verbal stimuli. In other words, reading the word rather than naming the color is the prepotent response. However, in the Stroop task, participants are given the instruction to ignore the word and instead read the color. We would argue that in this situation, the task instructions serve as context that allows the participant to inhibit the prepotent response tendency to read the word.

In many ways, one might argue that such a definition of context could be overly broad or inclusive, as almost any type of stimulus could be seen as forming a context representation. However, we would argue that context information or representations can be distinguished, both theoretically and empirically, from other related concepts in important ways. As an example, we argued that the representation and maintenance of context can be distinguished from both the constructs of short-term memory and working memory (Barch, Carter, & Cohen, 2003; Cohen, Barch, Carter, & Servan-Schreiber, 1999; Cohen & Servan-Schreiber, 1992). Short-term memory is often used to refer to processes necessary to temporarily

store recently presented information in a form somewhat close to the form in which it was presented (e.g., without conceptual transformation). Such information held in short-term memory may or may not have implications for how one should respond in a future situation. In contrast, our definition of context information is that it must be information that has relevance for later behavior, but it may not correspond to the actual identity of previously presented information. In fact, our assumption is that context representations are almost always transformations of the identity of specific stimuli into their meaning for behavior, which often may no longer include the initial identity of the stimulus. We do think context processing is much more closely aligned with conceptions of working memory, which is commonly defined as the collection of processes responsible for the on-line maintenance and manipulation of information necessary to perform a cognitive task (Baddeley, 1994). However, we do not believe that context processing and working memory are interchangeable constructs. Instead, we view context representations as a subset of representations within working memory, which govern how other representation held within either working memory or long-term memory are used.

An important insight that has emerged from our work is that the context processing functions of our model demonstrate how a single underlying mechanism, operating under different task conditions, might subserve three cognitive functions that are often treated as independent: attention (selection and support of task-relevant information for processing), active memory (on-line maintenance of such information), and inhibition (suppression of task-irrelevant information). When a task involves competing, task-irrelevant processes (as in the Stroop task), it is often assumed that a dedicated inhibitory function is responsible for suppressing, or overriding, these irrelevant processes. However, in our model, there is no dedicated mechanism for inhibition. Rather, context representations accomplish the same effect by providing top-down support for task-relevant processes, allowing these to compete effectively against irrelevant ones. In contrast, when a task involves a delay between a cue and a later contingent response, it is usually assumed that a working memory function is involved. Once again, there is no dedicated mechanism for this function in our model. Rather, the mechanism used to represent context information is used to maintain task relevant information against the interfering and cumulative effects of noise over time. Thus, both for tasks that tap "inhibition" and for those that tap "working memory," the same mechanism is involved; it is simply a matter of the behavioral conditions under which it operates (i.e., the source of interference) that lead us to label it as having an "inhibitory" or a "working memory" function. Furthermore, under both types of conditions, context representations serve an attentional function by selecting task-relevant information for processing over other potentially competing sources of information. Thus, in all circumstances, the

same context processing mechanism is involved. We hypothesize that this context processing mechanism is impaired in schizophrenia. Consequently, we suggest that disturbances in context processing may form a common basis for many of the deficits observed across multiple cognitive domains in schizophrenia, including attention, inhibition, working memory, and language processing.

In many ways, this notion of deficits in the representation of context is related to the notion of goal neglect discussed by West and Bowry in another chapter in this volume and in previous work (West, 1999; West & Alain, 2000). It is also related to similar ideas put forth by Duncan (Duncan, Emslie, Williams, Johnson, & Freer, 1996). For example, one can think of context information as representations of goals that are held in memory to guide and constrain ongoing processing. Difficulties in the sustained active maintenance of such representations can lead to a number of behavioral deficits on cognitive control tasks as described by West. As discussed in greater detail later, our model of context processing and the associated neurobiological mechanisms provides some hypotheses as to disturbances that might contribute to deficits in context processing and/or goal neglect.

Further, our hypotheses regarding the role of context processing in cognitive control (and the influence of such maintained context representations to overcome the negative influences of interference and conflict) are consistent with the role of "executive attention" in the working memory model put forth by Engle and colleagues (Engle & Kane, 2004). Specifically, Engle suggests the term executive attention refers to the ability to maintain even a single bit of information (e.g., a goal) in working memory in the face of a variety of sources of conflict. As with our theory, Engle and colleagues do not draw a strong line between mechanisms involved in the maintenance of information and those involved in the control and manipulation of ongoing processing. For example, Engle argues that although there may be multiple components of working memory capacity, the component most closely linked to success in a number of real-world outcomes (e.g., new vocabulary learning) is the ability to use attentional control to maintain goal-relevant information when there is interference or competition for other processes or stimuli. This is in contrast to the type of working memory model put forth by Baddeley, which does postulate a stricter segregation of storage/maintenance processes and central executive processes (Baddeley, 1996; Baddeley, 1986, 2000; Baddeley & Hitch, 1994), although more recently Baddeley emphasized the importance of attention control for working memory function (Baddeley, 1993; Baddeley & Logie, 1999). Thus, our model can be thought of as being highly compatible with that of Engle and colleagues. The primary differences between the models are ones of emphasis and the experimental domains upon which they have focused. Engle and colleagues have primarily focused on studies of healthy young adults, using both an individual differences approach and standard

cognitive psychology methodology. In contrast, our own model focused as heavily on special populations (i.e., schizophrenia, older adults) as on healthy young adults and integrated computational and cognitive neuroscience methodologies with cognitive experimental ones.

CONTEXT PROCESSING IN SCHIZOPHRENIA

In many ways, our hypotheses about context processing deficits in schizophrenia are very similar to earlier suggestions about the nature of cognitive deficits in this disorder put forth by researchers such as Shakow (Shakow, 1962). For example, Shakow suggested that "we see particularly the various difficulties created by context . . . It is as if, in the scanning process which takes place before the response to a stimulus is made, the schizophrenic is unable to select out the material relevant for optimal response (p. 4)" (Shakow, 1962). There is now a growing body of evidence from a variety of different task domains that supports the idea that individuals with schizophrenia have a deficit in the ability to represent and maintain context information, including tasks drawn from the domains of working memory, selective attention, inhibition, and language processing.

AX-CPT Task

One task that we used in numerous studies is a version of the classic Continuous Performance Test (Rosvold, Mirsky, Sarason, Bransome, & Beck, 1956) known as the AX-CPT (Cohen et al., 1999; Servan-Schreiber, Cohen, & Steingard, 1996). This test was specifically designed as a measure of context representation and maintenance. As shown in Figure 6.1a, in this task, participants are presented with cue-probe pairs and told to make a target response to an "X" (probe) but only when it follows an "A" (cue), and a nontarget response otherwise. A correct response to "X" depends upon maintaining the "context" provided by the cue ("A" or not-"A"). One change to the standard AX-CPT was to increase the frequency of target ("AX") trials so that they occur with a high frequency (70%, see Figure 6.1b), with the remaining 30% of trials distributed across three types of nontarget trials ("BX", "AY", and "BY" where "B" refers to any non-"A" cue and "Y" refers to any non-"X" probe). This creates two types of biases that can be used to probe the integrity of context processing. The first bias, or prepotent response, is that participants expect to make a target response when they see an "X" probe, because this is the correct response on most of the trials (87.5% of trials in which an X is presented). On "BX" trials, participants have to use the context provided by the "B" cue to inhibit this bias to respond target to an "X" (which would lead to a false alarm). Thus, impaired context representations will lead to poor performance on "BX" trials, because the context provided by the "B" cue would not be available

A

Target is an X following an A

B

| | | CUE | |
|-------|-----------|---------------------|-----------------------|
| | | VALID | INVALID |
| PROBE | TARGET | A-X (70%) | "B-X" (10%) |
| | NONTARGET | A-Y (10%) | "B-Y" (10%) |

to override the tendency to want to response target to the "X." The second bias is that participants expect to make a target response after they see an "A" cue, because most of the time an "X" follows the "A" cue (87.5% of "A" cue trials). However, on trials in which the "A" is not followed by an "X", this predictive aspect of context actually *creates* the tendency to false alarm. Thus, intact representations of context will hurt performance on "AY" trials, because context induces an invalid expectancy, leading to worse "AY" than "BX" performance. In contrast, individuals with impaired context representations should show worse "BX" than "AY" performance. In other words, individuals with impaired context representations (such as individuals with schizophrenia) should show worse "BX" performance, but *better* "AY" performance, than individuals with intact context representations.

A second manipulation included in the AX-CPT task is to examine maintenance of context information and the initial representation of context by manipulating the delay between the cue and probe (see Figure 6.1c). When the interval or delay between the cue and the probe is lengthened from, for example, 1 second to 5 or 10 seconds, context must be actively maintained within *working memory* (supported by prefrontal cortex in this and other theories of working memory function; see below). One prediction of

A number of prior studies found results with this AX-CPT task that provide evidence consistent with an impairment in context processing in schizophrenia (Barch et al., 2001; Barch et al., 2003; Cohen et al., 1999; Javitt, Shelley, Silipo, & Lieberman, 2000; Servan-Schreiber et al., 1996; Stratta, Daneluzzo, Bustini, Casacchia, & Rossi, 1998; Stratta, Daneluzzo, Bustini, Prosperini, & Rossi, 2000). For example, earlier work in our lab (Barch et al., 2001) demonstrated that as predicted, individuals with schizophrenia (first episode, neuroleptic naïve; see Figure 6.2a) made significantly more "BX" errors than did healthy controls, but significantly fewer "AY" errors, particularly at the longer delay. This pattern of both impaired and improved performance suggests that context representations are less available or less able to influence processing in individuals with schizophrenia, leading to an inability to override the incorrect tendency to respond target to the "X" on "BX" trials. In addition, such impairments in the processing of context lead to less predictive use of context on "AY" trials (which normally leads to errors on "AY" trials). In addition, the individuals with schizophrenia were significantly slower than controls on correct "BX" trials (suggesting increased interference from the "X" cue when there is a need to override a prepotent response tendency). However, the individuals with schizophrenia were not significantly slower than controls on "AY" trials, despite the fact that patients are typically overall slower on all tasks (see Figure 6.2b). This lack of an RT difference on AY trials again suggests that patients did not use context in a predictive fashion, leading them to actually experience less of a context-induced interference effect.

Across a number of different studies, results with the AX-CPT have consistently demonstrated that individuals with schizophrenia have deficits in the representation of context, in that they show the predicted patterns of increased BX errors and RTs, but not increased AY errors (or even decreased) and RTs. However, the answer to the question of whether patients with schizophrenia have a deficit in the maintenance of context




FIGURE 6.2. The graph on the left depicts "BX" and "AY" errors in the AX-CPT task in participants from (Barch, Carter et al., 2001); the graph on the right depicts "BX" and AY RTs from the same participants.

(e.g., effects exacerbated at the longer delay) and a deficit in the initial representation of context (e.g., deficits at short delay) is unclear. If patients with schizophrenia were primarily experiencing deficits in the representation of context, but not in maintenance, then performance deficits should be equal across the short and long delay. In contrast, if patients had deficits in representation and maintenance, then deficits should be worse at the long as compared to short delay. Some studies have shown that individuals with schizophrenia show an increase in context processing deficits at the long as compared to short delay (Cohen et al., 1999; Elvevag, Duncan, & McKenna, 2000; Javitt et al., 2000; Servan-Schreiber et al., 1996; Stratta et al., 1998; Stratta et al., 2000), whereas other studies have not (Barch et al., 2001; Barch et al., 2003; MacDonald et al., in submission; Perlstein, Dixit, Carter, Noll, & Cohen, 2003). However, close examination of the patterns of data across these different studies suggests that two different factors may be influencing whether context processing deficits are significantly greater at the long as compared to short delay in patients with schizophrenia as compared to controls. The first factor is whether the patients are early in the course of illness (e.g., first episode) or more chronic. To date, the studies with first episodic patients have not, for example, found significant increases in "BX" errors or RTs at the long as compared to the short delay (Barch et al., 2001; Barch et al., 2003; MacDonald et al., in submission) or have at least found smaller increases as a function of delay than in studies with chronic patients (Javitt et al., 2000). This suggests that early in the course of illness, patients with schizophrenia experience deficits in the initial representation of context but not further deficits in the maintenance of context. In contrast, studies with chronic patients do indicate that "BX" errors and RTs are significantly increased at the long as compared short delay (Cohen et al., 1999; Elvevag et al., 2000; Javitt et al., 2000; Servan-Schreiber et al., 1996; Stratta et al., 1998; Stratta et al., 2000), suggesting that chronic patients may have deficits in context maintenance and in initial context representation. A second factor is the performance of controls. In some studies with chronic patients, the controls also showed an increase in "BX" errors or RTs from the short to long delay, making it difficult to detect a differentially greater increase in patients (Cohen et al., 1999; Javitt et al., 2000). At this point, it is not entirely clear why some healthy controls show reductions in context processing at the long versus short delay, though it is possible that this may be related to some changes in the ability to represent and/or maintain context that occurs with age (see below for further discussion of aging and context processing) (Braver et al., 2001) or that can vary as a function of factors such as fluid intelligence (Burgess & Braver). Additional studies more directly examining stage of illness effects or longitudinal changes in context processing among both patients with schizophrenia and healthy controls will help to clarify the magnitude of deficits in




FIGURE 6.3. The graph on the left depicts "BX" and "AY" errors in the AX-CPT task in participants from (Barch et al., 2003); the graph on the right depicts "BX" and "AY" RTs from the same participants.

the initial representation of context versus the maintenance of context in schizophrenia.

In other work, we also examined the degree to which such context processing deficits are specific to individuals with schizophrenia as compared to other psychiatric disorders. For example, one study compared chronic individuals with schizophrenia to individuals with nonpsychotic major depression and found that unlike individuals with schizophrenia, individuals with major depression did not show any evidence of impairments in context processing (Cohen et al., 1999). In more recent work (Barch et al., 2003), we compared the specificity of context processing deficits to individuals with schizophrenia versus individuals with other psychotic disorders (e.g., major depression with psychotic features, bipolar disorder, delusional disorder, psychotic disorder NOS). In this study, participants were entered into the study upon their first contact with psychiatric services and tested before they were ever administered antipsychotic medications. Participants were then followed longitudinally, with repeat testing after four weeks and a confirmation of their diagnostic status at six months. At study admission, the individuals with psychotic disorders other than schizophrenia demonstrated a very similar pattern of deficits in context processing to that shown by individuals with schizophrenia. This included more "BX" errors and slower "BX" RTs as compared to controls, but the same or "AY" errors and no difference in "AY" RTs. Again, this suggests that both the individuals with schizophrenia and the individuals with other psychotic disorders were less able to use context representations to inhibit an incorrect target response on "BX" trials and less able to use context to predict potential targets on "AY" trials.

However, at four weeks, the individuals with other psychotic disorders, but not the individuals with schizophrenia, began to show improvements in context processing. As shown in Figure 6.3, by four weeks, the individuals with other psychotic disorders no longer differed from controls in "BX" performance (either errors or RT), but they did show significant performance differences when compared to individuals with schizophrenia. Taken together, we argued that such results suggest that context processing deficits are a more stable component of schizophrenia, perhaps forming part of the vulnerability to this disorder. In contrast, context processing deficits may be more state-related in other psychotic disorders (Barch et al., 2003).

Others research found that individuals at risk for schizophrenia, and who presumably share liability for this disorder, also display deficits in context processing on the AX-CPT task. For example, MacDonald and colleagues showed that nonpsychotic siblings of patients with schizophrenia show a similar performance pattern on the AX-CPT to that found with schizophrenia patients, demonstrating increased "BX" errors and RTs and decreased "AY" errors and no difference in "AY" RTs (MacDonald,




FIGURE 6.4. The graph on the left depicts "BX" and "AY" errors in the AX-CPT task in controls and individuals with schizotypal personality disorder (SPD) from (Barch, Mitropoulou et al., in press); the graph on the right depicts "BX" and "AY" RTs from the same participants.

Pogue-Geile, Johnson, & Carter, 2003). In our own work, we examined context processing deficits in individuals with schizotypal personality disorder, a disorder thought to share genetic liability with schizophrenia. As shown in Figure 6.4, we also found a pattern of deficits on the AX-CPT among individuals with schizotypal personality disorder that is indicative of a disturbance in context processing (Barch et al., in press). Like individuals with schizophrenia, the individuals with schizotypal personality disorder displayed increased "BX" errors, but not "AY" errors, combined with increased "BX" RTs, but not "AY" RTs. However, none of these effects were significantly exacerbated with delay in the individuals with schizotypal personality disorder. Taken together, these results suggest that individuals with schizotypal personality disorder also have difficulty utilizing contextual information to govern behavioral responding, but that this deficit is not worsened by increasing the delay over which such information must be maintained.

Other Measures of Context Processing

A number of studies using tasks other than the AX-CPT also provided evidence for deficits in context processing in schizophrenia. For example, a growing number of studies suggest that deficits in context processing can be observed in the domain of language comprehension and production. Very interesting evidence for context processing deficits in schizophrenia comes from a number of studies examining an ERP component called the N400, which is thought to index the degree of consistency or relatedness between a stimulus (e.g., word) and the preceding context (which could be a single word, a sentence, or an entire discourse; Brown & Hagoort, 1993; Kutas & Hillyard, 1980). The N400 is a negative going component of an ERP waveform that occurs approximately 400 msec after the onset of the stimulus of interest. In cognitively intact individuals, words that are not consistent with the preceding context elicit a larger N400 than words that are consistent with the preceding context.

A number of such N400 studies using different types of paradigms have provided evidence for context processing deficits in schizophrenia. For example, Condray used a lexical decision task in which pairs of words were presented that were either semantically related or semantically unrelated (Condray, Steinhauer, Cohen, van Kammen, & Kasperek, 1999; for related studies, see Grillon, Ameli, & Glzer, 1991). Controls demonstrated a significant priming effect in N400s (larger N400s to unrelated as compared to related words), whereas drug-free individuals with schizophrenia did not show an N400 priming effect. This suggests that patients were unable to use the context provided the first word to bias processing of the second word. Salisbury (Salisbury, O'Donnell, McCarley, Nestor, & Shenton, 2000) used a paradigm in which he presented four-word sentences to participants,

of the form "THE NOUN WAS ADJECTIVE/VERB." In this paradigm, the adjective or verb was always congruent with the noun. Controls showed significantly reduced N400s to the sentence final adjective/verbs as compared to the nouns presented earlier in the sentence, consistent with the idea that the controls use the noun to generate contextual representations that can help facilitate processing of consistent semantic information. However, the patients with schizophrenia did not show any reduction in N400 to the sentence final words as compared to the nouns, suggesting that patients were unable to either develop or use such contextual representations.

In more recent work, Sitnikova and colleagues (Sitnikova, Salisbury, Kuperberg, & Holcomb, 2002) examined N400s in a paradigm involving two clause sentences. The first clause ended with a homograph that had asymmetrical meanings (a dominant meaning and a subordinate meaning), and the second clause started with a strong semantic associate of the dominant meaning of the homograph. On half the trials, the first clause biased the dominant meaning of the homograph (so that the second clause should be consistent and not elicit an N400). On the other half of the trials, the first clause biased the subordinate meaning of the homograph (so that the second clause should be inconsistent and should elicit an N400). The control participants demonstrated reliably larger N400s in the inconsistent second-clause condition as compared to the consistent second-clause condition. This suggests that controls were able to use the context in the first clause to facilitate processing of the subordinate meaning of the homograph and suppress the dominant meaning. In contrast, the patients with schizophrenia showed a significantly smaller N400 than controls to the dominant meaning of the homograph when it followed a context that should have biased the subordinate meaning, and in fact showed no differences in N400 to consistent and inconsistent second clauses (for related studies, see (Adams et al., 1993; Niznikiewicz et al., 1997; Ohta, Uchiyama, Matsushima, & Toru, 1999; Olichney, Iragui, Kutas, Nowacki, & Jeste, 1997; Strandburg et al., 1997; Titone, Holzman, & Levy, 2002; Titone, Levy, & Holzman, 2000)). Again, such results suggest that the patients with schizophrenia were less able to use the context provided by the first part of the sentence to bias ongoing processing. Of note, we do not mean to imply that we think the N400 itself is necessarily generated by the same context representation and maintenance processes that support AX-CPT task performance. However, variations in the N400 (and the cognitive processes generating this ERP component) may be influenced by the integrity of such context representation mechanisms. If so, then examining variations in the N400 may allow one to make inference about context representation and maintenance processes that modulate the types of integration or semantic process that give rise to the N400.

Studies in the domain of selective attention have also been used as evidence for deficits in context processing. For example, a number of studies using the color-word Stroop task found that individuals with

schizophrenia are less able to inhibit the prepotent response to read the word, as evidenced by increased errors in the incongruent condition (color and word conflict) or by an increase in the total Stroop effect in RT (Barch et al., 1999; Barch, Carter, Hachten, & Cohen, 1999; Carter et al., 1992; Chen, Wong, Chen, & Au, 2001; Cohen et al., 1999; Elvevåg et al., 2000; Henik et al., 2002; Taylor, Kornblum, & Tandon, 1996). Further, in recent work, we used the process dissociation techniques developed by Jacoby to estimate the contributions of both word naming and color reading to Stroop performance in schizophrenia and found that color reading estimates are reduced whereas word reading estimates increased (Barch, Carter, & Cohen, in press). As described previously, our theory posits that task instructions to attend to color and ignore words serve as context representations in the Stroop tasks that normally allow one to inhibit the tendency to read words. Thus, we argued that Stroop task deficits in schizophrenia are consistent with a deficit in context processing.

Silverstein and colleagues suggested that deficits in context processing and contextual integration among individuals with schizophrenia extend even to the level of basic perceptual processing. (Silverstein et al., 2000). It is not yet clear whether the type of contextual processing measured in these paradigms is the same as that measured in tasks such as AX-CPT. However, such results raise the intriguing possibility that deficits in context processing account for deficits among individuals with schizophrenia both on high-level cognitive tasks and in more basic sensory and perceptual domains.

INTERRELATIONSHIP OF CONTEXT PROCESSING AND OTHER COGNITIVE CONSTRUCTS

As noted earlier, one of our hypotheses is that deficits in context processing serve as a common underlying impairment across a wide variety of cognitive domains, including working memory, inhibition/selective attention, and language processing. To test this hypothesis, we conducted a study in which we administered several different tasks to individuals with schizophrenia, along with healthy controls and depressed individuals (see Figure 6.5) (Cohen et al., 1999). We picked tasks that had traditionally been associated with different cognitive domains: (a) working memory with the AX-CPT; (b) inhibition/selective attention with the color-word Stroop; and (c) language processing with a lexical disambiguation task. However, each of the tasks included similar manipulations designed to increase their reliance on context representation and maintenance. As shown in Figure 6.5, each task included some type of contextual cue, a prepotency manipulation (either naturally occurring or induced in the task), and a delay manipulation (between the context and the need to use the context). The AX-CPT task was administered as described earlier. The Stroop task was modified to be a "switching" Stroop in which trials randomly

Context Tasks

In each task, there was:

- 1) a delay between the cue (context) and response
- 2) an asymmetry in the relative prepotency of competing responses

| Task | Context | Prepotency | Delay |
|------------------------|---|--|---|
| AX-CPT | "A" or "Not-A" cue | 70% Target (A-X) Sequences | 1 or 5 sec cue-probe interval |
| Stroop | Word Reading or Color Naming Instruction | Word Reading more prepotent than color naming | 1 or 5 sec cue-stimulus interval |
| Lexical Disambiguation | Sentence that biases a dominant or subordinate completion Example: race/rage ("ra_e") Race: Fred and Bob dashed to the finish line in a tie. Rage: Fred and Bob had bad tempers and often became quite angry. | One completion of an item ("ra_e") more frequent (race) in the absence of context than other completion (rage) | Sentence order, either: neutral -> Bias or bias -> Neutral |

FIGURE 6.5. Description of the tasks and manipulations of context representation and delay (Cohen et al., 1999).

alternated from color-naming to word-reading, with the type of trial indicated by a cue (i.e., context) that occurred at the start of each trial. Thus, participants needed to use the context provided by this cue to inhibit the tendency to read the word on color naming trials and needed to update this context information on a trial-by-trial basis. In addition, we manipulated the delay between the cue and the onset of the stimulus in order to assess maintenance of context. In the lexical disambiguation task, participants were presented with words missing one letter that could be completed in one of two ways, with one completion occurring more frequently than another (in the general population). These "missing letter" stimuli were preceded by sentences that could bias the participant toward the more dominant completion or toward the less frequent completion. As such, participants needed to use the context provided by the preceding sentences to modulate the tendency to complete the letter strings with the dominant versus subordinate completion. As with the other two tasks, we also manipulated the delay between this contextual information and the occurrence of the missing letter stimuli to assess context maintenance.

The results of this study indicated that, for the most part, individuals with schizophrenia were impaired across all three cognitive domains and tasks in the conditions that we would argue most strongly tap the need to represent and maintain context information. For example, the patients with schizophrenia again demonstrated increased "BX" errors as compared to

the controls, but no increased AY errors, as well as no increased "AX" errors, especially with a long delay. On the Stroop task, the patients with schizophrenia demonstrated significantly more errors than did the controls, particularly in the incongruent condition of the color naming task, the condition we argue is most dependent on context processing. On the lexical disambiguation task, the individuals with schizophrenia demonstrated a clear pattern indicative of impaired context processing, which included: (a) demonstrating the same asymmetry in production of the dominant versus subordinate completions as the controls in the absence of any biasing context; (b) the production of fewer subordinate completions, particularly with a long delay, when the sentence context should have biased such completions; and (c) the production of fewer dominant completions than controls when the sentence context should have biased them toward such completions (ruling out the possibility that patients simply had a dominant response bias). More importantly, however, were the results of the analyses examining cross-task relationships. We found that among the total sample and just within the individuals with schizophrenia, performance on each of the three tasks in conditions dependent on context processing were strongly and selectively interrelated, whereas performance on conditions not dependent on context (but psychometrically matched) were not related. Such results provide support for the hypothesis that deficits in a wide variety of cognitive domains in schizophrenia may result from disturbances in contextual processing.

In other studies, we also found that performance on tasks specifically designed to measure context processing, such as the AX-CPT, are strongly correlated with performance on tasks measuring other cognitive processes, which we would argue are also dependent on context processing. For example, in a large sample of healthy younger and older adults, we found that performance on the AX-CPT was correlated with performance on a number of standard measures of working memory, including the reading span, a highly demanding digit span, and an "N-back" task (Keys, Barch, Braver, & Janowsky, submitted). In the study described previously with individuals who have schizotypal personality disorder, we found that AX-CPT performance was strongly correlated with performance on an "N-Back" task, which measures working memory, and on an Eriksson Flanker task, which measures selective attention (Barch et al., in press). Again, such findings are consistent with the hypothesis that deficits in context processing may contribute to disturbances in a variety of related cognitive domains.

THE NEUROBIOLOGY OF CONTEXT PROCESSING

We hypothesized that the representation and maintenance of context processing are subserved by a specific set of neural mechanisms. In

particular, we postulate that representations of context information are housed within the dorsolateral portion of the prefrontal cortex (DL-PFC) and actively maintained there when task demands require such active maintenance (O'Reilly, Braver, & Cohen, 1999). Further, we hypothesize that the dopamine (DA) projections to DL-PFC regulate the access to such context information, insulating this information from the interfering effects of noise over intervals in which the information must be sustained, while at the same time allowing for the appropriate updating of such context information when needed (Braver & Cohen, 2000). These hypotheses are consistent with the broader neuroscience literature, in which active maintenance in the service of control is a commonly ascribed function to PFC (Fuster, 1989; Goldman-Rakic, 1987; Miller & Cohen, 2001), and the DA system is widely held to modulate the active maintenance properties of PFC (Cohen, Braver, & Brown, 2002; Luciana, Collins, & Depue, 1998; Sawaguchi, Matsumura, & Kubota, 1990; Williams & Goldman-Rakic, 1995). In our model, the context processing functions of cognitive control critically depend upon DL-PFC and DA system interactions. As a consequence, the model predicts that individuals and populations with impairments in either or both DL-PFC or the DA system should demonstrate specific patterns of impaired cognitive control related to the processing of context.

ROLE OF PREFRONTAL CORTEX

As noted above, a large number of functional neuroimaging studies have demonstrated that prefrontal cortex is activated when individuals have to maintain information in working memory (Cabeza & Nyberg, 2000). However, the majority of these studies used tasks that may or may not tap context processing, and they were not specifically designed to selectively measure context processing. However, we and our colleagues conducted a number of neuroimaging studies using tasks, such as the AX-CPT and the switching Stroop, that were specifically designed to measure context processing. For example, our early research with the AX-CPT demonstrated that dorsolateral PFC and ventrolateral PFC showed selective increases in activity when context representations had to be maintained over a long delay as compared to short delay (Barch et al., 1997; Braver & Cohen, 2001). In contrast, these same PFC regions did not show increased activity when the AX-CPT task was made more difficult by degrading the stimuli (but not changing maintenance demands). This suggests that dorsolateral PFC regions are not simply activated whenever the task gets more challenging, although the anterior cingulate did respond to this difficulty manipulation, leading to subsequent research on the role of the anterior cingulate in conflict detection (Botvinick et al., 2001). Such findings dissociating the control processes subserved by the dorsolateral PFC relative to other

brain regions (such as the anterior cingulate) have been replicated by other studies as well (MacDonald, Cohen, Stenger, & Carter, 2000). Of interest, the only brain regions to show a sustained increase in activity when required to maintain context information were the dorsolateral and ventrolateral PFC regions. This is in contrast to the patterns of brain activity often found during working memory tasks, which is typically much more widespread, including regions such as the parietal cortex, frontomedial motor regions, and other subcortical regions (e.g., cerebellum, basal ganglia). We argued that this selective responsivity of PFC regions to the context maintenance manipulation provides strong support for their central role in supporting this particular component of cognitive control.

We also examined how challenging context processing in healthy individuals alters task-related PFC activity. For example, we developed a version of the AX-CPT in which participants are presented with distractor items during the delay between the cue and the probe. Our hypothesis was that if deficits on the AX-CPT among individuals with schizophrenia were due to disturbances in the ability to maintain context information, disturbing the ability to maintain context information in healthy adults should elicit the same types of task deficits as found in individuals with schizophrenia. Consistent with this hypothesis, we found that adding distractors during the delay between the cue and the probe in the long-delay conditions of the AX-CPT increased the number of "BX" errors (and slowed "BX" RTs) that healthy participants made, but *decreased* the number of "AY" errors, consistent with a reduced ability to use context information. In a subsequent functional neuroimaging study using the same task design, we found that the addition of distractions and impairments in task performance was accompanied by a selective decrease in dorsolateral PFC activity but no change or even an increase in ventrolateral PFC activity (Braver & Cohen, 2001). The results of this study and others have provided some clues about the different contributions that dorsolateral versus ventrolateral PFC play in context processing. More specifically, we argued that ventrolateral regions of PFC may serve a more general role in phonological processing or rehearsal that may in no way be selective or specific to context representations, an idea put forth by many other researchers as well. In contrast, it may be that dorsolateral PFC (particularly in the left hemisphere) may be more specifically involved in the development and/or maintenance of context representations.

We also examined the patterns of brain activity found in individuals with schizophrenia during the performance of tasks such as the AX-CPT. In one such study, medication naïve first episode individuals with schizophrenia and controls were presented with trials that had either a short- or a long-delay period between the cue and the probe (Barch et al., 2001). The trials were designed so that four images of the brain were acquired during each trial, allowing us to glean some rough information about the time course




FIGURE 6.6. Illustration of task-related brain activity in dorsolateral PFC during the AX-CPT. The image displays a coronal slice through the brain 25 mm anterior to the anterior commissure plane. The graph illustrated percent fMRI signal change during each scan of the trial, relative to the first scan of the short delay condition, for each group in each condition.

of brain activity during context maintenance. As shown in Figure 6.6, the healthy controls once again demonstrated a sustained increase in dorsolateral PFC activity during the long as compared to short-delay condition. However, the medication naïve individuals with schizophrenia did not show any significant increased in dorsolateral PFC activity in response to the context maintenance demand. In contrast, as shown in Figure 6.7, the individuals with schizophrenia did show increased delay related activity in ventrolateral PFC regions, activity that did not differ in magnitude from controls. Again, this provides evidence for the fact that dorsolateral and ventrolateral regions of PFC play different roles in context processing and are differentially impaired in schizophrenia.

Such findings of impaired dorsolateral PFC activity during the performance of context processing tasks have now been replicated several times, in chronic and in first episode patients (MacDonald, in press; Perlstein, Dixit, Carter, Noll, & Cohen, 2003). Further, recent research suggests that such dorsolateral PFC deficits are specific to individuals with schizophrenia as compared to individuals with other psychotic disorders, who show a different pattern of potentially increased dorsolateral PFC activity (MacDonald et al., in submission).




FIGURE 6.7. Illustration of task-related brain activity in ventrolateral PFC during the AX-CPT. The image displays a coronal slice through the brain 10 mm anterior to the anterior commissure plane. The graphs for both the left and right ventrolateral regions display the increase in task-related brain activity from the short- to the long-delay condition.

ROLE OF DOPAMINE

As described above, our models also postulate a central role for dopamine in context processing and hypothesize that impaired dopamine function in dorsolateral PFC contributes to context processing deficits. There is a large body of evidence in nonhuman primates and humans suggesting that dopamine plays an important role in working memory more generally (Arnsten, Cai, Murphy, & Goldman-Rakic, 1994; Barch, in press; Brozoski, Brown, Rosvold, & Goldman, 1979; Cai & Arnsten, 1997; Castner, Williams, & Goldman-Rakic, 2000; de Sonneville, Nijokiktjen, & Bos, 1994; Goldman-Rakic, 1995; Kimberg & D'Esposito, 2003; Kimberg, D'Esposito, & Farah, 1997; Luciana & Collins, 1997; Luciana, Collins, & Depue, 1995; Luciana et al., 1998; Luciana, Depue, Arbisi, & Leon, 1992; Mattay et al., 1996; Mattay et al., 2000; Mattay et al., 2003; Mehta et al., 2000; Mehta, Swainson, Gogilvie, Sahakian, & Robbins, 2001; Mintzer & Griffiths, 2003; Muller, von Cramon, & Pollmann, 1998; Sawaguchi & Goldman-Rakic, 1994; Williams & Goldman-Rakic, 1995). This includes evidence that non-specific dopamine agonists, such as amphetamine, can improve working memory in medicated patients with schizophrenia (Barch & Carter, in preparation; Daniel et al., 1991; Goldberg, Bigelow, Weinberger, Daniel, & Kleinman, 1991) and that changes in D1 receptor availability in DLPFC are associated with working memory impairment in schizophrenia (Abi-Dargham et al., 2002). Unfortunately, however, there is little direct evidence yet for a role for dopamine specifically in context processing in humans, as compared to working memory more generally. However, we have recently completed a double-blind placebo controlled study examining the influence of d-amphetamine on AX-CPT performance in healthy individuals.

In this study we used both our standard version of the AX-CPT and the "interference" version described previously, in which distractors appeared between the cue and the probe. As noted earlier, including such distractors normally elicits a pattern of performance in controls analogous to that found in schizophrenia (e.g., increased "BX" errors and RT; decreased "AY" errors). Participants were 12 healthy controls who participated in two experimental sessions spaced no more than one week apart. In each session, participants performed both the baseline and interference conditions. At each session, participants were orally administered either placebo or D-AMP, at a dosage of 0.25 mg/kg of body weight, at the beginning of the session. The drug on each day was administered in a double-blind manner (with an unlabeled opaque capsule), and drug order was randomly counterbalanced across participants. Because D-AMP effects peak after one to two hours, and are stable over the following one to two hours (Angrist, Corwin, Barlett, & Cooper, 1987) participants were tested approximately two hours post-ingestion. To analyze the data, we examined a measure of context sensitivity that we refer to as d' -context, a signal-to-noise measure comparing hits to "AX" trials and false alarms to "BX" trials. As shown in Figure 6.8, we found that under placebo, the addition of distractors during the cue-probe period of long-delay trials reduced sensitivity to context, as indexed by a reduced d' -context (e.g., increased "AX" misses and "BX" false alarms). However, this effect was significantly reduced with D-amphetamine. More specifically, there was a significant main effect of D-amphetamine on d' -context ($F(1,11) = 6.41, p < .05$) and a significant drug \times condition interaction ($F(1,11) = 6.31, p < .05$). Planned contrasts revealed that under placebo, context sensitivity was reduced in the interference condition ($F(1,11) = 18.39, p < .01$), but that there was no significant difference between the two conditions under D-AMP ($F(1,11) = 0.32, p > .10$) (see Figure 6.8). Results of studies such as this provide some more specific evidence for a relationship between dopamine function and context processing, at least in healthy individuals.

On a related note, the finding that dopaminergic agents can influence context processing may help to shed light on the mechanisms that by which stress influences working memory. In another chapter in this book, Sliwinski and colleagues review evidence that higher perceived levels of stress are associated with cognitive impairment, particularly impaired working memory, on both a between-person basis (i.e., comparing individuals with high and low levels of perceived stress) and a within-person basis (i.e., comparing times with high and low perceived stress within the same person). Sliwinski and colleagues put forth the hypothesis that this relationship may reflect that fact that individuals coping with a stressor may use cognitive resources to suppress stress-related thoughts, leaving fewer resources available for other cognitive tasks. However, stress is also known to alter dopamine function, at least acutely, and some animal research has shown a

d' -Context Under Placebo and D-Amphetamine




FIGURE 6.8. Graph illustrating change in d' -context as a function of adding interfering distractors, both under placebo and under d-amphetamine.

detrimental impact of stress on working memory via dopaminergic mechanisms in prefrontal cortex (Arnsten & Goldman-Rakic, 1998). As such, it is possible that stress influences working memory, and potentially context processing, via disruption of the dopamine system. This hypothesis is not meant to be an alternative to the one put forth by Sliwinski, but rather may provide a different level of explanation for the same mechanism.

RELATIONSHIP OF CONTEXT PROCESSING TO CLINICAL SYMPTOMS IN SCHIZOPHRENIA

One important question that arises when discussing context processing deficits in schizophrenia is whether such disturbances are present in all individuals with schizophrenia with equal severity, or whether the severity of such context processing deficits are related to the severity of specific symptoms. We and others found that context processing deficits in schizophrenia appear to be strongly related to the severity of a constellation of behaviors referred to as disorganization symptoms, which include disorganized behavior (dressing in an unusual manner, behaving oddly in public), attentional problems, and difficulties in producing goal directed speech (e.g., loose associations (Barch et al., 1999; Barch & Carter, 1998; Barch

al., 2003; Cohen et al., 1999; MacDonald et al., in submission; Perlstein, Carter, Noll, & Cohen, 2001; Perlstein et al., 2003; Stratta et al., 2000). Such findings are interesting in light of research suggesting that representations of context information are important for guiding coherent ongoing language production (Levelt, 1989). Further, such associations are consistent with the research reviewed above, suggesting that impairments in language comprehension, homograph interpretation/meaning selection, and semantic priming may be related to impairments in the ability to use or maintain contextual information.

More recently, we also found some evidence that deficits in context processing may also be related to disturbances in the ability to regulate the influence of emotion on cognitive processing in schizophrenia. More specifically, we found that individuals with impaired context processing show greater language production disruptions (in the form of unclear references) when discussing affectively negative as compared to affective neutral topics (Burbridge & Barch, 2002). There is a growing literature in affective neuroscience examining the role that different cognitive control functions play in emotional regulation and the role that dorsolateral PFC plays in emotional regulation (Gray, 2001; Gray, Braver, & Raichle, 2002; Ochsner, Bunge, Gross, & Gabrieli, 2002). One hypothesis is that intact representations of context are also important for the regulation of emotion because such information provides clues as to what emotional responses are appropriate in given situations and a means by which to alter emotional responses that are not compatible with current task demands. As such, it will be interesting in future research to examine the role that the representation and maintenance of context plays in successful emotional regulation and the potential influence of disordered context processing on disturbed emotional functioning in disorders such as schizophrenia.

CONTEXT PROCESSING IN OTHER POPULATIONS

Individuals with schizophrenia are not the only population thought to experience changes in prefrontal function that may be associated with changes in dopamine function. For example, individuals with Parkinson's disease are known to have disturbances in dopamine function and disturbances in tasks designed to measure working memory and other aspects of cognitive control (Zgaljardic, Borod, Foldi, & Mattis, 2003). In addition, individuals with attention deficit hyperactivity disorder are thought to suffer from disturbances in dopamine function and prefrontal function and again show deficits on a range of cognitive control tasks (Casey et al., 1997; Aron, Dowson, Sahakian, & Robbins, 2003; Nigg, 2003). Of interest, healthy older adults represent a nonpathological population that also experiences cognitive control problems that may be related to changes in prefrontal and dopamine function. For example, a growing body of work suggests

that healthy aging involves systematic changes in prefrontal function as measured by functional imaging techniques (Cabeza, 2001; Grady, 1998) and involves changes in cognitive functions thought to be associated with prefrontal function (Balota, Dolan, & Ducheck, 2000; Jacoby, Debner, & Hay, 2001), both of which may be associated with changes in dopamine function with aging (Arnsten et al., 1994; Arnsten, Cai, Steere, & Goldman-Rakic, 1995; de Keyser, De Backer, Vauquelin, & Ebinger, 1990; Goldman-Rakic & Brown, 1981; Suhara et al., 1991; Volkow et al., 1998)). Further, in many ways, the types of cognitive deficits shown by healthy older individuals are very similar to those found in individuals with schizophrenia. The chapter by West and Bowry in this volume nicely articulates many of the types of cognitive deficits shown by healthy older adults, which are similar to deficits seen in individuals with schizophrenia. For example, both individuals with schizophrenia and healthy older adults show increased incongruent errors in the Stroop and an increased total Stroop effect in RT (Spieler, Balota, & Faust, 1996; Verhaeghen & De Meersman, 1998; West & Baylis, 1998), and both populations have more difficulties with color naming in a switching Stroop that varies the task to be performed on a trial-by-trial basis, as compared to blocked trials of color naming (Cohen et al., 1999; West, in press). Similar to our interpretations of the source of these deficits in schizophrenia, West attributed such Stroop task deficit in healthy older adults to disturbances in context processing.

We directly studied context processing and related functions in healthy aging (e.g., working memory, inhibition) and also found evidence that healthy older adults show deficits in context processing (Braver et al., 2001), which are strongly correlated with deficits in measures of working memory and inhibition (Keys et al., submitted). For example, in recent work, we found that healthy young-old adults (ages 65 to 75) show deficits in context representation, in the form of increased "BX" RTs as compared to younger adults, but not increased "AY" RTs (Braver, Satpute, Keys, & Racine, in press). Further, healthy much older adults (76+) demonstrated additional deficits in context maintenance, in that their "BX" worsened as a function of delay, while "AY" performance improved.

RELATIONSHIPS AMONG ANTERIOR CINGULATE, DOPAMINE FUNCTION, AND COGNITIVE CONTROL

In addition, West and others examined additional components of cognitive control in healthy aging, such as error processing as indexed by an ERP component referred to as the error related negativity (ERN). The ERN is thought to index either error monitoring and/or correction (Falkenstein, Hohnsbein, Hoorman, & Blanke, 1991; Falkenstein, Hoorman, Christ, & Hohnsbein, 2000; Gehring, Goss, Coles, Meyer, & Donchin, 1993) or conflict processing (Botvinick et al., 2001), critical components of cognitive

control. A number of researchers believe that one of the main generators of the ERN is the anterior cingulate, and that alterations in the ERN may also reflect changes in dopamine function in the anterior cingulate (Botvinick et al., 2001; Holroyd & Coles, 2002). As reviewed by West, the amplitude of the ERN is reduced in healthy older adults across a wide range of task paradigms (Band & Kok, 2000; Falkenstein, Hoorman, & Hohnbein, 2001; Nieuwenhuis et al., 2002; West, *in press*), a finding consistent with the research on reduced dopamine function in healthy aging. Similarly, a growing number of studies also suggest reduced ERNs in individuals with schizophrenia, even when there are no behavioral differences in conditions thought to be associated with increased conflict (Alain, McNeely, Yu, Christensen, & West, 2002; Bates, Kiehl, Laurens, & Liddle, 2002; Carter, MacDonald III, Ross, & Stenger, 2001; Kopp & Rist, 1999) and decreased anterior cingulate activity, a potential generator of the ERN (Carter et al., 2001). Again, such results in schizophrenia may reflect abnormalities in dopamine function, which influences anterior cingulate and dorsolateral PFC function. Such results suggest that both individuals with schizophrenia and healthy older adults may experience deficits in error monitoring/conflict processing and deficits in context processing.

Despite the many similarities in the profiles of cognitive disturbances shown by individuals with schizophrenia and healthy older adults, there are also intriguing differences. For example, in studies of context processing in healthy older adults using the AX-CPT task, deficits in at least the young-older adults are apparent primarily in RTs rather than errors. In contrast, individuals with schizophrenia manifest context-processing deficits in both errors and RTs on the AX-CPT. In the Stroop task, research examining process-dissociation estimates of color naming and word reading has found increased word reading estimates in older adults, but no changes in color naming estimates (Spieler et al., 1996). In contrast, in schizophrenia, we have found both increased word reading and decreased color naming estimates (Barch et al., *in press*). In addition, as noted by West, studies of healthy aging populations have not found increased correct trial related negativities in ERP studies (referred to as the CRN; Band & Kok, 2000; Falkenstein et al., 2001; West, *in press*), whereas studies of schizophrenia have reported such increased CRNs (Alain et al., 2002; Mathalon et al., 2002). One argument is that increased CRNs reflect additional conflict because of inadequate goal representations or context representations that normally serve to reduce conflict in information processing. These differences in the performance of individuals with schizophrenia and healthy older adults suggest that although these two populations may share some common cognitive and neurobiological disturbances, there are also important differences. For example, one possibility is that the severity of context processing deficits is simply worse in individuals with schizophrenia as compared to healthy adults. Such a hypothesis would be consistent with

the fact that individuals with schizophrenia show problems in both errors and RTs and alterations in both color and word processing estimates (alterations in only word processing estimates among older adults might reflect a less severe disturbance). However, it is somewhat less clear why a less severe context processing deficit would lead to reductions only in the ERN and not the CRN, as compared to the concurrent reductions in the ERN and the CRN found in schizophrenia. Future research more directly comparing the performance of individuals with schizophrenia and healthy older adults in the same exact paradigms may help clarify some of these issues and better delineate the similarities and differences in the profiles of cognitive disturbances found in these two populations.

It is not necessarily surprising that both individuals with schizophrenia and healthy older individuals show similar patterns of cognitive deficits if both populations experience changes in prefrontal function associated with dopamine changes. Although the source of the prefrontal cortex and dopamine changes in schizophrenia and healthy aging may be very different, it may still be the case that a final common pathway of changes in prefrontal function leads to somewhat similar cognitive deficits in schizophrenia and healthy aging. However, at the same time, it is also clear that there are many differences between individuals with schizophrenia and healthy older adults, both in the severity of their cognitive deficits and the associated symptoms and phenomenology found in schizophrenia. Such differences may reflect several factors. First, many researchers believe that at least some aspects of the neurobiological pathology found in schizophrenia are neurodevelopmental in origin and are thus present throughout the life span. In contrast, changes in prefrontal function in healthy older adults may begin to occur much later in life, with intact prefrontal function early in life. The eventual outcome of neurodevelopmental changes in prefrontal function may be very different than changes that occur only later in life, because such prefrontal changes could then influence the course of learning, skill acquisition, and cognitive development in individuals with schizophrenia, which may have additional contributions to the profile of cognitive disturbances found in schizophrenia. Second, it is unlikely that changes in prefrontal cortex and dopamine function are the only neurobiological disturbances found in schizophrenia. As such, the interaction of multiple sources of neurobiological and cognitive abnormalities in schizophrenia may contribute to many of the cognitive deficits/symptoms found in schizophrenia that are not present in healthy aging (although healthy aging also may involve changes in brain regions other than prefrontal cortex).

SUMMARY

In this chapter, we described our theory regarding one component of cognitive control, the representation and maintenance of context, that we argue is

important for the ongoing control of thoughts, behaviors, and emotion. We argued that this particular component of cognitive control is supported by dopamine function in dorsolateral prefrontal cortex. As described in more detail above, this theory regarding cognitive control is highly consistent with the theories put forth by a number of authors of other chapters in this volume, including the ideas put forth by Engle, West, Sliwinski and their colleagues. Further, we argued that populations that experience disturbances in either or both dopamine function and dorsolateral prefrontal cortex function experience deficits in the processing of context, and we reviewed empirical data supporting this assertion. One important area for future research will be to better delineate the similarities and difference in the profiles of cognitive and neurobiological disturbances across populations thought to suffer from disturbances in prefrontal and dopamine function. By better understanding the relationships between specific types of cognitive control deficits and particular kinds of neurobiological disturbances, we can better validate, modify, and/or expand theories regarding the neurobiology of cognitive control. We have also suggested that the processes involved in the representation and maintenance of context may play a critical role in emotion regulation, though considerably more empirical and theoretical work is needed to examine this hypothesis.

Author Note

Research was supported by grants from the National Alliance for Research on Schizophrenia and Depression, the Dana Foundation, and the National Institute of Mental Health.

References

- Abi-Dargham, A., Mawlawi, O., Lombardo, I., Gil, R., Martinez, D., Huang, Y., et al. (2002). Prefrontal dopamine D1 receptors and working memory in schizophrenia. *Journal of Neuroscience*, 22, 3708-3719.
- Adams, J., Fauz, S. F., Nestor, P. G., Shenton, M. E., Marcy, B., Smith, S. R., et al. (1993). ERP abnormalities during semantic processing in schizophrenia. *Schizophrenia Research*, 10, 247-257.
- Alain, C., McNeely, H. E., Yu, H., Christensen, B. K., & West, R. (2002). Neurophysiological evidence of error monitoring deficits in patients with schizophrenia. *Cerebral Cortex*, 12, 840-846.
- Angrist, B., Corwin, J., Barlett, B., & Cooper, T. (1987). Early pharmacokinetics and clinical effects of oral d-amphetamine in normal subjects. *Biological Psychiatry*, 22, 1357-1368.
- Arnsten, A. F., Cai, J. X., Murphy, B. L., & Goldman-Rakic, P. S. (1994). Dopamine D1 receptor mechanisms in the cognitive performance of young adult and aged monkeys. *Psychopharmacology*, 116, 143-151.
- Arnsten, A. F., Cai, J. X., Steere, J. C., & Goldman-Rakic, P. S. (1995). Dopamine D2 receptor mechanisms contribute to age-related cognitive decline: the effects of quinpirole on memory and motor function in monkeys. *Journal of Neuroscience*, 15, 3429-3439.
- Arnsten, A. F., & Goldman-Rakic, P. S. (1998). Noise stress impairs prefrontal cortical cognitive function in monkeys: Evidence for a hyperdopaminergic mechanism. *Archives of General Psychiatry*, 55, 362-368.
- Aron, A. R., Dowson, J. H., Sahakian, B. J., & Robbins, T. W. (2003). Methylphenidate improves response inhibition in adults with attention-deficit/hyperactivity disorder. *Biological Psychiatry*, 54, 1465-1468.
- Baddeley, A. (1994). The magical number seven: Still magic after all these years. *Psychological Review*, 101, 353-356.
- Baddeley, A. (1996). Exploring the central executive. *Quarterly Journal of Experimental Psychology*, 49, 5-28.
- Baddeley, A. D. (1986). *Working memory*. New York: Oxford University Press.
- Baddeley, A. D. (1993). Working memory or working attention? In A. D. Baddeley & L. Weiskrantz (Eds.), *Attention: Selection, awareness, and control: A tribute to Donald Broadbent* (pp. 152-170). Oxford: Clarendon Press.
- Baddeley, A. D. (2000). The episodic buffer: A new component of working memory? *Trends in Cognitive Science*, 4, 417-423.
- Baddeley, A. D., & Hitch, G. J. (1994). Developments in the concept of working memory. *Neuropsychology*, 8, 485-493.
- Baddeley, A. D., & Logie, R. H. (1999). Working memory: The multiple-component model. In A. Miyake & P. Shah (Eds.), *Models of working memory: Mechanisms of active maintenance and executive control* (pp. 28-61). Cambridge: Cambridge University Press.
- Balota, D. A., Dolan, P. O., & Duchek, J. M. (2000). Memory changes in healthy older adults. In E. Tulving & F. I. M. Craik (Eds.), *The Oxford handbook of memory* (pp. 395-409). New York: Oxford University Press.
- Band, G., & Kok, A. (2000). Age effects on response monitoring in a mental rotation task. *Biological Psychology*, 51, 201-221.
- Barch, D. M. (in press). Pharmacological manipulations of human working memory. *Psychopharmacology*.
- Barch, D. M., Braver, T. S., Akbudak, E., Conturo, T., Ollinger, J., & Snyder, A. V. (2001). Anterior cingulate cortex and response conflict: Effects of response modality and processing domain. *Cerebral Cortex*, 11, 837-848.
- Barch, D. M., Braver, T. S., Nystrom, L. E., Forman, S. D., Noll, D. C., & Cohen, J. D. (1997). Dissociating working memory from task difficulty in human prefrontal cortex. *Neuropsychologia*, 35, 1373-1380.
- Barch, D. M., & Carter, C. S. (1998). Selective attention in schizophrenia: Relationship to verbal working memory. *Schizophrenia Research*, 33, 53-61.
- Barch, D. M., & Carter, C. S. (in preparation). *The influence of d-amphetamine on cognition and language in medicated patients with schizophrenia*. Manuscript in preparation.
- Barch, D. M., Carter, C. S., Braver, T. S., McDonald, A., Sabb, F. W., Noll, D. C., et al. (2001). Selective deficits in prefrontal cortex regions in medication naive schizophrenia patients. *Archives of General Psychiatry*, 50, 280-288.

- rch, D. M., Carter, C. S., & Cohen, J. D. (2003). Context processing deficit in schizophrenia: Diagnostic specificity, 4-week course, and relationships to clinical symptoms. *Journal of Abnormal Psychology*, 112, 132-143.
- rch, D. M., Carter, C. S., & Cohen, J. D. (in press). Process dissociation analyses of Stroop performance in schizophrenia. *Neuropsychology*.
- rch, D. M., Carter, C. S., Hachten, P. C., & Cohen (1999). The "benefits" of distractibility: The mechanisms underlying increased Stroop effects in schizophrenia. *Schizophrenia Bulletin*, 24, 749-762.
- rch, D. M., Carter, C., Perlstein, W., Baird, J., Cohen, J., & Schooler, N. (1999). Increased Stroop facilitation effects in schizophrenia are not due to increased automatic spreading activation. *Schizophrenia Research*, 39, 51-64.
- rch, D. M., Mitropoulou, V., Harvey, P. D., New, A. S., Silverman, J. M., & Siever, L. J. (in press). Context processing deficits in schizotypal personality disorder. *Journal of Abnormal Psychology*.
- ates, A. T., Kiehl, K. A., Laurens, K. R., & Liddle, P. F. (2002). Error-related negativity and correct response negativity in schizophrenia. *Clinical Neurophysiology*, 113, 1454-1463.
- otvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. C. (2001). Conflict monitoring and cognitive control. *Psychological Review*, 108, 624-652.
- raver, T. S. (1997). *Mechanisms of cognitive control: A neurocomputational model*. Unpublished doctoral dissertation, Carnegie Mellon University, Pittsburgh, PA.
- raver, T. S., Barch, D. M., & Cohen, J. D. (1999a). Cognition and control in schizophrenia: A computational model of dopamine and prefrontal function. *Biological Psychiatry*, 46, 312-328.
- raver, T. S., Barch, D. M., & Cohen, J. D. (1999b). *Mechanisms of cognitive control: Active memory, inhibition, and the prefrontal cortex* (Technical Report No. PDP.CNS.99.1). Pittsburgh PA: Carnegie Mellon University.
- raver, T. S., Barch, D. M., Keys, B. A., Carter, C. S., Cohen, J. D., Kaye, J. A., et al. (2001). Context processing in older adults: Evidence for a theory relating cognitive control to neurobiology in healthy aging. *Journal of Experimental Psychology: General*, 130, 746-763.
- raver, T. S., & Bongiolatti, S. R. (2002). The role of the frontopolar prefrontal cortex in subgoal processing during working memory. *NeuroImage*, 15, 523-536.
- raver, T. S., & Cohen, J. D. (1999). Dopamine, cognitive control, and schizophrenia: The gating model. *Progress in Brain Research*, 121, 327-349.
- raver, T. S., & Cohen, J. D. (2000). On the control of control: The role of dopamine in regulating prefrontal function and working memory. In S. Monsell & J. Driver (Eds.), *Attention and performance XVIII* (pp. 713-738). Cambridge, MA: MIT Press.
- raver, T. S., & Cohen, J. D. (2001). Working memory, cognitive control, and the prefrontal cortex: Computational and empirical studies. *Cognitive Processing*, 2, 25-55.
- raver, T. S., Cohen, J. D., & Barch, D. M. (2002). The role of the prefrontal cortex in normal and disordered cognitive control: A cognitive neuroscience perspective. In D. T. Stuss & R. T. Knight (Eds.), *Principles of Frontal Lobe Function* (pp. 428-448). Oxford: Oxford University Press.
- Braver, T. S., Cohen, J. D., & McClelland, J. L. (1997). *An integrated computational model of dopamine function in reinforcement learning and working memory*. Paper presented at the Society for Neuroscience Abstracts, New Orleans.
- Braver, T. S., Cohen, J. D., & Servan-Schreiber, D. (1995). Neural network simulations of schizophrenic performance in a variant of the CPT-AX: A predicted double dissociation. *Schizophrenia Research*, 15, 110.
- Braver, T. S., Satpute, A. B., Keys, B. A., & Racine, C. A. (in press). Context processing and context maintenance in healthy aging and early-stage dementia of the Alzheimer's type. *Psychology and Aging*.
- Brown, C., & Hagoort, P. (1993). The processing nature of the N400: evidence from masked priming. *Journal of Cognitive Neuroscience*, 5, 34-44.
- Brozoski, T. J., Brown, R. M., Rosvold, H. E., & Goldman, P. S. (1979). Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science*, 205, 929-931.
- Burbridge, J., & Barch, D. M. (2002). Emotional valence and reference disturbance in schizophrenia. *Journal of Abnormal Psychology*, 111, 186-191.
- Burgess, G. C., & Braver, T. S. (submitted). *Proactive interference effects on working memory can be modulated by expectancy: Evidence for dual mechanisms of cognitive control*. Manuscript submitted for publication.
- Cabeza, R. (2001). Functional neuroimaging of cognitive aging. In R. Cabeza & A. Kingstone (Eds.), *Handbook of functional neuroimaging of cognition*. Cambridge, MA: MIT Press.
- Cabeza, R., & Nyberg, L. (2000). Imaging cognition II: An empirical review of 275 PET and fMRI studies. *Journal of Cognitive Neuroscience*, 12, 1-47.
- Cai, J. X., & Arnsten, A. F. T. (1997). Dose-dependent effects of the dopamine D1 receptor agonists A77636 or SKF81297 on spatial working memory in aged monkeys. *The journal of pharmacology and experimental therapeutics*, 283, 183-189.
- Carter, C. S., MacDonald III, A. W., Ross, L. L., & Stenger, V. A. (2001). Anterior cingulate cortex activity and impaired self-monitoring of performance in patients with schizophrenia: an event-related fMRI study. *American Journal of Psychiatry*, 158, 1423-1428.
- Carter, C. S., Robertson, L. C., & Nordahl, T. E. (1992). Abnormal processing of irrelevant information in schizophrenia: Selective enhancement of Stroop facilitation. *Psychiatry Research*, 41, 137-146.
- Casey, B. J., Castellanos, F. X., Giedd, J. N., Marsh, W. L., Hamburger, S. D., Schubert, A. B., et al. (1997). Implication of right frontostriatal circuitry in response inhibition in an attention-deficit/hyperactivity disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36.
- Castner, S. A., Williams, G. V., & Goldman-Rakic, P. S. (2000). Reversal of antipsychotic induced working memory deficits by short term dopamine D1 receptor stimulation. *Science*, 287, 2020-2022.
- Chapman, J., & McGhie, A. (1962). A comparative study of disordered attention in schizophrenia. In *Depression and personality* (pp. 487-500). Missing info
- Chen, E. Y. H., Wong, A. W. S., Chen, R. Y. L., & Au, J. W. Y. (2001). Stroop interference and facilitation effects in first-episode schizophrenia patients. *Schizophrenia Research*, 48, 29-44.

- Cohen, J. D., Barch, D. M., Carter, C., & Servan-Schreiber, D. (1999). Context-processing deficits in schizophrenia: Converging evidence from three theoretically motivated cognitive tasks. *Journal of Abnormal Psychology*, 108, 120-133.
- Cohen, J. D., Braver, T. S., & Brown, J. W. (2002). Computational perspectives on dopamine function in prefrontal cortex. *Current Opinion in Neurobiology*, 12, 223-229.
- Cohen, J. D., & Servan-Schreiber, D. (1992). Context, cortex and dopamine: A connectionist approach to behavior and biology in schizophrenia. *Psychological Review*, 99, 45-77.
- Condray, R., Steinhauer, S. R., Cohen, J. D., van Kammen, D. P., & Kasperek, A. (1999). Modulation of language processing in schizophrenia: effects of context and haloperidol on the event-related potential. *Biological Psychiatry*, 45, 1336-1355.
- Cornblatt, B. A., & Keilp, J. G. (1994). Impaired attention, genetics, and the pathophysiology of schizophrenia. *Schizophrenia Bulletin*, 20, 31-62.
- Daniel, D. G., Weinberger, D. R., Jones, D. W., Zigur, J. R., Coppola, R., Handel, S., et al. (1991). The effect of amphetamine on regional cerebral blood flow during cognitive activation in schizophrenia. *The Journal of Neuroscience*, 11, 1907-1917.
- de Keyser, J., De Backer, J.-P., Vauquelin, G., & Ebinger, G. (1990). The effect of aging on the D1 dopamine receptors in human frontal cortex. *Brain Research*, 528, 308-310.
- de Sonneville, L. M., Nijokiktijen, C., & Bos, H. (1994). Methylphenidate and information processing: Part 1: Differentiation between responders and non-responders; Part 2: Efficacy in responders. *Journal of Clinical and Experimental Neuropsychology*, 16, 877-897.
- Duncan, J., Emslie, H., Williams, P., Johnson, R., & Freer, C. (1996). Intelligence and the frontal lobe: The organization of goal-directed behavior. *Cognitive Psychology*, 30, 257-303.
- Elvevåg, B., Duncan, J., & McKenna, P. J. (2000). The use of cognitive context in schizophrenia: An investigation. *Psychological Medicine*, 30, 885-897.
- Engle, R. W., & Kane, M. J. (2004). Executive attention, working memory capacity and a two-factor theory of cognitive control. In B. Ross (Ed.), *The psychology of learning and motivation* (Vol. 44). New York: Elsevier.
- Falkenstein, M., Hohnsbein, J., Hoorman, J., & Blanke, L. (1991). Effects of cross-modal divided attention on late ERP components: II. Error processing in choice reaction tasks. *Electroencephalography and Clinical Neurophysiology*, 78, 447-455.
- Falkenstein, M., Hoorman, J., Christ, S., & Hohnsbein, J. (2000). ERP components on reaction errors and their functional significance: a tutorial. *Biological Psychology*, 51, 87-107.
- Falkenstein, M., Hoorman, J., & Hohnsbein, J. (2001). Changes in error-related ERPs with age. *Experimental Brain Research*, 138, 258-262.
- Fuster, J. M. (1989). *The prefrontal cortex* (2nd ed.). New York: Raven Press.
- Gehring, W. J., Goss, B., Coles, M. G. H., Meyer, D. E., & Donchin, E. (1993). A neural system for error detection and compensation. *Psychological Science*, 4, 385-390.
- Gold, J. M., Carpenter, C., Randolph, C., Goldberg, T. E., & Weinberger, D. R. (1997). Auditory working memory and Wisconsin Card Sorting Test performance in schizophrenia. *Archives of General Psychiatry*, 54, 159-165.
- Gold, J. M., Randolph, C., Carpenter, C. J., Goldberg, T. E., & Weinberger, D. R. (1992). Forms of memory failure in schizophrenia. *Journal of Abnormal Psychology*, 101, 487-494.
- Goldberg, T. E., Bigelow, L. B., Weinberger, D. R., Daniel, D. G., & Kleinman, J. E. (1991). Cognitive and behavioral effects of the coadministration of dextroamphetamine and haloperidol in schizophrenia. *American Journal of Psychiatry*, 148, 78-84.
- Goldman-Rakic, P. S. (1987). Circuitry of primate prefrontal cortex and regulation of behavior by representational memory. In F. Plum & V. Mountcastle (Eds.), *Handbook of Physiology - The Nervous System V* (Vol. 5, pp. 373-417). Bethesda, MD: American Physiological Society.
- Goldman-Rakic, P. S. (1995). Cellular basis of working memory. *Neuron*, 14, 477-485.
- Goldman-Rakic, P. S., & Brown, R. M. (1981). Regional changes of monoamines in cerebral cortex and subcortical structures of aging rhesus monkeys. *Neuroscience*, 6, 177-187.
- Grady, C. L. (1998). Brain imaging and age-related changes in cognition. *Experimental Gerontology*, 33, 661-673.
- Gray, J. R. (2001). Emotional modulation of cognitive control: Approach-withdrawal states double-dissociate spatial from verbal two-back task performance. *Journal of Experimental Psychology: General*, 130, 436-452.
- Gray, J. R., Braver, T. S., & Raichle, M. E. (2002). Integration of emotion and cognition in the lateral prefrontal cortex. *Proceedings of the National Academy of Sciences USA*, 99, 4115-4120.
- Green, M. F., Kern, R. S., Braff, D. L., & Mintz, J. (2000). Neurocognitive deficits and functional outcome in schizophrenia: are we measuring the "right stuff"? *Schizophrenia Bulletin Special Issue: Psychosocial treatment for schizophrenia*, 26, 119-136.
- Grillon, C., Ameli, R., & Glzer, W. M. (1991). N400 and semantic categorization in schizophrenia. *Biological Psychiatry*, 29, 467-480.
- Henik, A., Carter, C. S., Salo, R., Chaderjian, M., Kraft, L., Nordahl, T. E., et al. (2002). Attentional control and word inhibition in schizophrenia. *Psychiatry Research*, 110, 137-149.
- Holroyd, C. B., & Coles, M. G. (2002). The neural basis of human error processing: reinforcement learning, dopamine, and the error-related negativity. *Psychological Review*, 109, 679-709.
- Jacoby, L. L., Debnar, J. A., & Hay, J. F. (2001). Proactive interference, accessibility bias, and process dissociations: Valid subjective reports of memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 27, 686-700.
- Javitt, D. C., Shelley, A., Silipo, G., & Lieberman, J. A. (2000). Deficits in auditory and visual context-dependent processing in schizophrenia. *Archives of General Psychiatry*, 57, 1131-1137.
- Keys, B. A., Barch, D. M., Braver, T. S., & Janowsky, J. S. (submitted). Task sensitivity to age differences in working memory: Relative superiority of the N-back paradigm. Manuscript submitted for publication.
- Kimberg, D. Y., & D'Esposito, M. (2003). Cognitive effects of the dopamine receptor agonist pergolide. *Neuropsychologia*, 41, 1020-1027.

- Kimberg, D. Y., D'Esposito, M., & Farah, M. J. (1997). Effects of bromocriptine on human subjects depend on working memory capacity. *Neuroreport*, 8, 381-385.
- Kopp, B., & Rist, F. (1999). An event-related brain potential substrate of disturbed response monitoring in paranoid schizophrenia patients. *Journal of Abnormal Psychology*, 108, 337-346.
- Kutas, M., & Hillyard, S. A. (1980). Reading senseless sentences: Brain potentials reflect semantic incongruity. *Science*, 207, 203-205.
- Levelt, W. J. M. (1989). *Speaking: From intention to articulation*. Cambridge, Mass: MIT Press.
- Luciana, M., & Collins, P. F. (1997). Dopamine modulates working memory for spatial but not object cues in normal humans. *Journal of Cognitive Neuroscience*, 4, 58-68.
- Luciana, M., Collins, P. F., & Depue, R. A. (1995). *DA and 5-HT influences on spatial working memory functions of prefrontal cortex*. Paper presented at the Cognitive Neuroscience Society Second Annual Meeting, San Francisco, CA.
- Luciana, M., Collins, P. F., & Depue, R. A. (1998). Opposing roles for dopamine and serotonin in the modulation of human spatial working memory functions. *Cerebral Cortex*, 8, 218-226.
- Luciana, M., Depue, R. A., Arbisi, P., & Leon, A. (1992). Facilitation of working memory in humans by a D₂ dopamine receptor agonist. *Journal of Cognitive Neuroscience*, 4, 58-68.
- MacDonald, A., Carter, C. S., Kerns, J. G., Ursu, S., Barch, D. M., Holmes, A. J., et al. (in submission). *Specificity of prefrontal dysfunction and context processing deficits to schizophrenia in a never medicated first-episode psychotic sample*. Manuscript submitted for publication.
- MacDonald, A. W. (in press). Event-related fMRI study of context processing in dorsolateral prefrontal cortex of patients with schizophrenia. *Journal of Abnormal Psychology*.
- MacDonald, A. W., Cohen, J. D., Stenger, V. A., & Carter, C. S. (2000). Dissociating the role of the dorsolateral prefrontal cortex and anterior cingulate cortex in cognitive control. *Science*, 288, 1835-1838.
- MacDonald, A. W., Pogue-Geile, M. F., Johnson, M. K., & Carter, C. S. (2003). A specific deficit in context processing in the unaffected siblings of patients with schizophrenia. *Archives of General Psychiatry*, 60, 57-65.
- Mathalon, D. H., Dedor, M., Faustman, W. O., Gray, M., Askari, N., & Ford, J. M. (2002). Response-monitoring dysfunction in schizophrenia: An event-related brain potential study. *Journal of Abnormal Psychology*, 111, 22-41.
- Mattay, V. S., Berman, K. F., Ostrem, J. L., Esposito, G., Van Horn, J. D., Bigelow, L. B., et al. (1996). Dextroamphetamine enhances "neural network-specific" physiological signals: A positron-emission tomography rCBF study. *Journal of Neuroscience*, 15(August), 4816-4822.
- Mattay, V. S., Callicott, J. H., Bertolino, A., Heaton, I., Frank, J. A., Coppola, R., et al. (2000). Effects of dextroamphetamine on cognitive performance and cortical activation. *Neuroimage*, 12, 268-275.
- Mattay, V. S., Goldberg, T. E., Fera, F., Hariri, A. R., Tessitore, A., Egan, M. F., et al. (2003). Catechol O-methyltransferase val158-met genotype and individual variation in the brain response to amphetamine. *Proceedings of the National Academy of Sciences*, 100, 6186-6191.

- McClelland, J. L. (1993). Toward a theory of information processing in graded, random, and interactive networks. In D. E. Meyer & S. Kornblum (Eds.), *Attention and Performance XIV: Synergies in experimental psychology, artificial intelligence, and cognitive neuroscience* (pp. 655-688). Cambridge, MA: MIT Press.
- Mehta, M. A., Owen, A. M., Sahakian, B. J., Mavaddat, N., Pickard, J. D., & Robbins, T. W. (2000). Methylphenidate enhances working memory by modulating discrete frontal and parietal lobe regions in the human brain. *The Journal of Neuroscience*, 20, 1-6.
- Mehta, M. A., Swainson, R., Gogilvie, A. D., Sahakian, B. J., & Robbins, T. W. (2001). Improved short-term spatial memory but impaired reversal learning following the dopamine D₂ agonist bromocriptine in human volunteers. *Psychopharmacology*, 159, 10-20.
- Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, 21, 167-202.
- Mintzer, M., & Griffiths, R. R. (2003). Triazolam-amphetamine interaction: dissociation of effects of memory versus arousal. *Journal of Pharmacology*, 17, 17-29.
- Muller, U., von Cramon, Y., & Pollmann, S. (1998). D₁- versus D₂-receptor modulation of visuospatial working memory in humans. *The Journal of Neuroscience*, 18, 2720-2728.
- Nieuwenhuis, S., Ridderinkhof, K. R., Talsma, D., Coles, M. G. H., Hoyroyd, C. B., Koki, A., et al. (2002). A computational account of altered error processing in older age: Dopamine and error-related negativity. *Cognitive, Affective and Behavioral Neuroscience*, 2, 19-36.
- Nigg, J. T. (2003). Response inhibition and disruptive behaviors: toward a multi-process conception of etiological heterogeneity for ADHD combined type and conduct disorder early-onset type. *Annals of the New York Academy of Sciences*, 1008, 170-182.
- Niznikiewicz, M. A., O'Donnell, B. F., Nestor, P. G., Smith, L., Law, S., Karapellou, M., et al. (1997). ERP assessment of visual and auditory language processing in schizophrenia. *Journal of Abnormal Psychology*, 106, 85-94.
- Nuechterlein, K. H., & Dawson, M. E. (1984). Information processing and attentional functioning in the developmental course of schizophrenia disorders. *Schizophrenia Bulletin*, 10, 160-203.
- O'Reilly, R. C., Braver, T. S., & Cohen, J. D. (1999). A biologically-based computational model of working memory. In A. Miyake & P. Shah (Eds.), *Models of Working Memory: Mechanisms of Active Maintenance and Executive Control*. New York: Cambridge University Press.
- O'Reilly, R. C., Noelle, D. C., Braver, T. S., & Cohen, J. D. (2002). Prefrontal cortex and dynamic categorization tasks: representational organization and neuromodulatory control. *Cerebral Cortex*, 12, 246-257.
- Ochsner, K. N., Bunge, S. A., Gross, J. J., & Gabrieli, J. D. (2002). Rethinking feelings: an fMRI study of the cognitive regulation of emotion. *Journal of Cognitive Neuroscience*, 14, 1215-1229.
- Ohta, K., Uchiyama, M., Matsushima, E., & Toru, M. (1999). An event-related potential study in schizophrenia using Japanese sentences. *Schizophrenia Research*, 40, 159-170.

- Olichney, J. M., Iragui, V. J., Kutas, M., Nowacki, R., & Jeste, D. V. (1997). N400 abnormalities in late life schizophrenia and related psychoses. *Biological Psychiatry*, 42, 13-23.
- Park, S., & Holzman, P. S. (1992). Schizophrenics show spatial working memory deficits. *Archives of General Psychiatry*, 49, 975-982.
- Perlstein, W. H., Carter, C. S., Noll, D. C., & Cohen, J. D. (2001). Relation of prefrontal cortex dysfunction to working memory and symptoms in schizophrenia. *American Journal of Psychiatry*, 158, 1105-1113.
- Perlstein, W. M., Dixit, N. K., Carter, C. S., Noll, D. C., & Cohen, J. D. (2003). Prefrontal cortex dysfunction mediates deficits in working memory and prepotent responding in schizophrenia. *Biological Psychiatry*, 53, 25-38.
- Rosvold, H. E., Mirsky, A. F., Sarason, I., Bransome, E. D. Jr., Beck, L. H. (1956). A continuous performance test of brain damage. *Journal of Consulting Psychology*, 20, 343-350.
- Rumelhart, D. E., & McClelland, J. L. (1986). *Parallel distributed processing: Explorations in the microstructure of cognition* (Vols. 1 and 2). Cambridge, MA: MIT Press.
- Salisbury, D., O'Donnell, B. F., McCarley, R. W., Nestor, P. G., & Shenton, M. E. (2000). Event-related potentials elicited during a context-free homograph task in normal versus schizophrenic subjects. *Psychophysiology*, 37, 456-463.
- Sawaguchi, T., & Goldman-Rakic, P. S. (1994). The role of D1-dopamine receptor in working memory: Local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys performing an oculomotor delayed-response task. *Journal of Neurophysiology*, 71, 515-528.
- Sawaguchi, T., Matsumura, M., & Kubota, K. (1990). Effects of dopamine antagonists on neuronal activity related to a delayed response task in monkey prefrontal cortex. *Journal of Neurophysiology*, 63, 1401-1410.
- Servan-Schreiber, D., Cohen, J. D., & Steingard, S. (1996). Schizophrenic deficits in the processing of context: A test of a theoretical model. *Archives of General Psychiatry*, 53(Dec), 1105-1113.
- Shakow, D. (1962). Segmental set: A theory of the formal psychological deficit in schizophrenia. *Archives of General Psychiatry*, 6, 1-17.
- Silverstein, S. M., Kovacs, I., Corry, R., & Valone, C. (2000). Perceptual organization, the disorganization syndrome, and context processing in chronic schizophrenia. *Schizophrenia Research*, 43, 11-20.
- Sitnikova, T., Salisbury, D. F., Kuperberg, G., & Holcomb, P. J. (2002). Electrophysiological insights into language processing in schizophrenia. *Psychophysiology*, 39, 851-860.
- Spieler, D. H., Balota, D. A., & Faust, M. E. (1996). Stroop performance in healthy younger and older adults and in individuals with dementia of the Alzheimer's type. *Journal of Experimental Psychology: Human Perception and Performance*, 22, 461-479.
- Strandburg, R. J., Marsh, J. T., Brown, W. S., Asarnow, R. F., Guthrie, D., Harper, R., et al. (1997). Event-related potential correlates of linguistic information processing in schizophrenics. *Biological Psychiatry*, 42, 596-608.
- Stratta, P., Daneluzzo, E., Bustini, M., Casacchia, M., & Rossi, A. (1998). Schizophrenic deficits in the processing of context. *Archives of General Psychiatry*, 55, 186-187.

- Stratta, P., Daneluzzo, E., Bustini, M., Prosperini, P., & Rossi, A. (2000). Processing of context information in schizophrenia: relation to clinical symptoms and WCST performance. *Schizophrenia Research*, 44, 57-67.
- Suhara, T., Fukuda, H., Inoue, O., Itoh, T., Suzuki, K., Yamasaki, T., et al. (1991). Age-related changes in human D1 dopamine receptors measured by positron emission tomography. *Psychopharmacology*, 103, 41-45.
- Taylor, S. F., Kornblum, S., & Tandon, R. (1996). Facilitation and interference of selective attention in schizophrenia. *Journal of Psychiatric Research*, 30, 251-259.
- Titone, D., Holzman, P. S., & Levy, D. L. (2002). Idiom processing in schizophrenia: literal implausibility saves the day for idiom priming. *Journal of Abnormal Psychology*, 111, 313-320.
- Titone, D., Levy, D. L., & Holzman, P. S. (2000). Contextual insensitivity in schizophrenic language processing: evidence from lexical ambiguity. *Journal of Abnormal Psychology*, 109, 761-767.
- Verhaeghen, P., & De Meersman, L. (1998). Aging and the Stroop effect: a meta-analysis. *Psychology and Aging*, 13, 120-126.
- Volkow, N. D., Gur, R. C., Wang, G.-J., Fowler, J. S., Moberg, P. J., Ding, Y.-S., et al. (1998). Association between decline in brain dopamine activity with age and cognitive and motor impairment in healthy individuals. *American Journal of Psychiatry*, 155, 344-349.
- Weinberger, D. R., Berman, K. F., & Zec, R. F. (1986). Physiological dysfunction of dorsolateral prefrontal cortex in schizophrenia: I. Regional cerebral blood flow evidence. *Archives of General Psychiatry*, 43, 114-125.
- West, R. (1999). Age differences in lapses of intention in the Stroop task. *Journal of Gerontology: Psychological Sciences*, 54B, P34-P43.
- West, R. (in press). The effect of aging on controlled attention and conflict processing in the Stroop task. *Journal of Cognitive Neuroscience*.
- West, R., & Alain, C. (2000). Effects of task context and fluctuations of attention on neural activity supporting performance of the Stroop task. *Brain Research*, 873, 102-111.
- West, R., & Baylis, G. C. (1998). Effects of increased response dominance and contextual disintegration on the Stroop interference effect in older adults. *Psychology & Aging*, 13, 206-217.
- Williams, G. V., & Goldman-Rakic, P. S. (1995). Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature*, 376 (August), 572-575.
- Zgaljardic, D. J., Borod, J. C., Foldi, N. S., & Mattis, P. (2003). A review of the cognitive and behavioral sequelae of Parkinson's disease: relationship to frontostriatal circuitry. *Cognitive and Behavioral Neurology*, 16, 193-210.