

Neuroinflammation in the Amygdala Is Associated With Recent Depressive Symptoms

Wei Zhang, Jerrel Rutlin, Sarah A. Eisenstein, Yong Wang, Deanna M. Barch, Tamara Hershey, Ryan Bogdan, and Janine D. Bijstervosch

ABSTRACT

BACKGROUND: Converging evidence suggests that elevated inflammation may contribute to depression. Yet, the link between peripheral inflammation and neuroinflammation in depression is unclear. Here, using data from the UK Biobank, we estimated associations among depression, C-reactive protein (CRP) as a measure of peripheral inflammation, and neuroinflammation as indexed by diffusion basis spectral imaging–based restricted fraction (DBSI-RF).

METHODS: DBSI-RF was derived from diffusion-weighted imaging data ($N = 11,512$) for whole-brain gray matter (global-RF), and regions of interest in the bilateral amygdala (amygdala-RF) and hippocampus (hippocampus-RF), and CRP was estimated from blood (serum) samples. Self-reported recent depression symptoms were measured using a 4-item assessment. Linear regressions were used to estimate associations between CRP and DBSI-RFs with depression while adjusting for the following covariates: age, sex, body mass index, smoking, drinking, and medical conditions.

RESULTS: Elevated CRP was associated with higher depression symptoms ($\beta = 0.04$, false discovery rate–corrected $p < .005$) and reduced global-RF ($\beta = -0.03$, false discovery rate–corrected $p < .001$). Higher amygdala-RF was associated with elevated depression—an effect resilient to added covariates and CRP ($\beta = 0.02$, false discovery rate–corrected $p < .05$). Interestingly, this association was stronger in individuals with a lifetime history of depression ($\beta = 0.07$, $p < .005$) than in those without ($\beta = 0.03$, $p < .05$). Associations between global-RF or hippocampus-RF with depression were not significant, and no DBSI-RF indices indirectly linked CRP with depression (i.e., mediation effect).

CONCLUSIONS: Peripheral inflammation and DBSI-RF neuroinflammation in the amygdala are independently associated with depression, consistent with animal studies suggesting distinct pathways of peripheral inflammation and neuroinflammation in the pathophysiology of depression and with investigations highlighting the role of the amygdala in stress-induced inflammation and depression.

<https://doi.org/10.1016/j.bpsc.2023.04.011>

Major depressive disorder (MDD) is a worldwide health issue with profound emotional, cognitive, and economic consequences (1,2). Building upon meta-analytic studies that elevated inflammation is associated with depression (3,4), convergent evidence supports the theory that inflammation is a potential etiological factor contributing to the development of depression (5–14). For example, immunological challenges (e.g., influenza vaccine in humans, lipopolysaccharide administration in nonhuman animals, interferon treatment in both) induce depressive symptoms (15,16), whereas anti-inflammatory treatment reduces depressive symptoms (17–19). Furthermore, inflammation is prospectively associated with future depression, even when considering depression at baseline (4,13), and baseline depression has been prospectively associated with future elevations in inflammation (20), suggestive of a bidirectional relationship that may contribute to the widespread chronicity and recurrence among individuals with depression (3). Although a multitude of mechanisms may contribute to the relationship between peripheral inflammation

and depression, the putative depressogenic effects of inflammation are often attributed to neuroinflammation (21). However, there are limited studies of neuroinflammation in humans, and the available evidence is mixed due to small sample sizes in positron emission tomography studies on depression (22–24). While studies of microglial activity using translocator protein will be informative (25), increasing evidence suggests that biological correlates of complex behavioral phenotypes such as depression are characterized by small effects (26), highlighting the need to extend neuroinflammatory research into larger datasets. Here, we leverage the large-scale UK Biobank (UKB) dataset (analytic $n = 11,501$) to examine the relationship between neuroinflammation index and depressive symptoms.

Although neuroinflammation cannot be directly measured in humans in vivo, currently available neuroimaging methods including positron emission tomography, single-photon emission computed tomography, and magnetic resonance imaging (MRI) have been used for indirect assessment. Recent evidence from positron emission tomography and functional MRI

techniques indicates that putative neuroinflammation in depression might be most relevant for subcortical regions that subserve reward, motivation, and emotion processes. For example, increased plasma C-reactive protein (CRP) concentration was associated with decreased functional connectivity between the amygdala and ventromedial prefrontal cortex in patients with depression comorbid with anxiety disorder relative to healthy control participants (27), and depressive symptom severity was correlated with significant increases in translocator protein total density in the hippocampus in patients compared with healthy control participants (28). Interestingly, alterations in structure and function of the hippocampus and amygdala are also the most prominent neuroimaging markers for major depression disorder (29–32). These 2 subcortical regions are involved in a wide range of cognitive and emotional processes including emotion regulation, memory, and stress, and impairment in these functions are often evident in individuals with depression (33–35). It remains understudied, however, whether inflammation in these regions may contribute to depressive symptom severity and whether disruptions in these regions may potentially mediate the impact of peripheral inflammatory response on depression symptomatology. In this study, we used diffusion basis spectrum imaging (DBSI), a novel noninvasive technique, to estimate putative neuroinflammation in these regions of interest from diffusion-weighted imaging (DWI) data.

The DBSI approach is an extension of standard diffusion tensor imaging (DTI) that models the entire DWI signal as a linear combination of multiple anisotropic diffusion tensors and a spectrum of isotropic diffusion components (36,37). In comparison to DTI that describes water movement parallel and perpendicular to axon tracts, the DBSI approach has the advantage to improve specificity and sensitivity of diffusion properties by resolving multiple-tensor water diffusion resulting from axon injury, demyelination, and inflammation (36). In particular, DBSI-based restricted fraction (DBSI-RF), the metric of interest in this study, provides a putative indicator of neuroinflammation-related cellularity (i.e., the state of having cells), where higher values indicate increased immune cell infiltration relative to lower values (36). DBSI-RF has shown robust correlations with cell nuclei counts in brain tissues in a mouse model of autoimmune encephalomyelitis, with stain-quantitated cellularity from postmortem human central nervous system tissues and has been found to be greater in individuals with inflammation-related diseases such as multiple sclerosis, obesity, HIV, and Alzheimer's disease than in control participants (38–42). These studies suggest that DBSI-RF can be used as a proxy measure for inflammation-related cellularity in the brain.

For this study, our goal was to assess putative inflammation-related cellularity (i.e., DBSI-RF), specifically in the hippocampus, amygdala, and whole-brain gray matter in relation to depression. We further explored whether putative neuroinflammation mediates the association between peripheral inflammation and depression.

METHODS AND MATERIALS

Participants

The UKB is a large-scale study ($N > 500,000$ participants) designed to examine the genetic, environmental, biological,

and behavioral correlates of broad-spectrum health outcomes and related phenotypes (43). For the present study, we considered data from 16,182 participants who completed the baseline session (i.e., the first assessment center visit between 2006 and 2010), initial neuroimaging acquisition (i.e., the third assessment center visit from 2014 and 2019), and a web-based assessment session (2016–2017) of the UKB study.

Data were excluded for the following reasons: 1) mismatch between self-reported and genetic sex ($n = 130$), 2) missing data on a variable included in this study ($n = 1834$), 3) marked elevation of CRP levels (i.e., above 10 g/L; $n = 398$) that may reflect acute infection, injury, or disease (44), and 4) diseases associated with systemic inflammation (e.g., known inflammatory disease, autoimmune disease, HIV, hepatitis B or C; $n = 2319$). Thus, our final analytic sample consisted of 11,501 participants (Table 1). This study was conducted under the UKB Application ID 47267.

In a subset of sample for our secondary analysis on lifetime MDD phenotype, data of participants who met lifetime MDD criterion were excluded if they had reported other mood disorders (e.g., bipolar, schizophrenia) or were on treatment or medication for antipsychotics, whereas data of participants who did not meet lifetime MDD criterion were excluded if they had a history of depression or any mental problems, or on antidepressants (see detailed exclusion criterion in Table S2). The resulting sample had 6960 participants, including 1138 participants with lifetime MDD (Table 1).

Depression

The primary measure of depression used in our study was the total score of recent depressive symptoms (RDSs), which was completed on the same day as the neuroimaging session. This score was a sum of 4 items (scored on a 1–4 scale where 1 = not at all, 4 = nearly every day) assessing the presence of the following self-reported depressive symptoms over the past 2 weeks: 1) depressed mood, 2) unenthusiasm/disinterest, 3) tenseness/restlessness, and 4) tiredness/lethargy. The resulting sum score ranged between 4 and 16, where higher scores indicate more frequent and severe depressive symptoms. The RDS score has been previously validated against several commonly used depression scales, including the 9-item Patient Health Questionnaire (45). Variable ID numbers of these items in the UKB Data Showcase are summarized in Table S1.

In secondary analyses, we focused on a subset of participants of lifetime MDD phenotype (46). Lifetime MDD is defined based on the short form of the Composite International Diagnostic Interview (47) that was administered as part of the online mental health questionnaire (46). Briefly, participants were asked to identify their experiences regarding 2 core depression symptoms: 1) depressive mood and 2) loss of interest, for a period of 2 or more weeks. If “Yes” was the response to any of these questions, participants continued to indicate the lifetime number of depressive symptoms and their experiences on the other DSM-IV MDD symptoms during the worst episode: 3) feelings of worthlessness, 4) tiredness, 5) difficulty concentrating, 6) suicidal thoughts, 7) changes in sleeping pattern, and 8) changes in weight. This provides a 0 to 8 sum score of depression symptoms over 8 symptom questions, where higher scores indicate greater endorsement of depression

Table 1. Characteristics of Study Sample

	Full	Lifetime MDD		
		Yes	No	All
Sample Size				
Female, <i>n</i> (%)	6097 (53%)	1138 (67%)	2368 (45%)	3506 (50%)
Total, <i>N</i>	11,501	1693	5267	6960
Age, Years	62.41 (7.41)	61.17 (7.15)	63.05 (7.43)	62.59 (7.41)
Race, White, %	98.37%	99.17%	97.80%	98.13%
Body Mass Index	24.81 (2.74)	24.75 (2.81)	24.82 (2.70)	24.80 (2.73)
Serum C-Reactive Protein, mg/L	1.42 (1.46)	1.43 (1.45)	1.40 (1.44)	1.41 (1.44)
RDSs	5.15 (1.68)	5.68 ^a (2.02)	4.73 ^a (1.33)	4.97 (1.58)
DBSI-RF				
Gray matter	0.071 (0.011)	0.071 (0.011)	0.071 (0.011)	0.071 (0.011)
Hippocampus	0.037 (0.005)	0.037 (0.005)	0.037 (0.005)	0.037 (0.005)
Amygdala	0.028 (0.006)	0.028 (0.005)	0.028 (0.006)	0.028 (0.006)

Values are presented as mean (SD) except where indicated.

DBSI-RF, diffusion basis spectral imaging–based restricted fraction; MDD, major depressive disorder; RDS, recent depressive symptoms.

^aIndicates significant differences ($p < .00001$) in RDSs between the phenotype and the full sample such that individuals with lifetime MDD had a higher RDS score and individuals without lifetime MDD had a lower RDS score than that of the full study sample.

symptoms. Participants were also asked to indicate the frequency and duration of the core symptoms, as well as their psychological impairment (i.e., whether they interfere with their roles, life, and activities). An individual participant is considered to have lifetime MDD when their sum score ≥ 5 and experienced symptoms “almost every day” or “every day,” with a duration of “most of the day” or “all day,” and symptoms impaired psychosocial functioning “somewhat” or “a lot” (46). Following previous studies (48,49), we further applied exclusion criterion to ensure phenotype specificity to depression (e.g., excluding participants reporting bipolar disorder or schizophrenia). Details are summarized in Table S2. As shown in Table 1, the mean depression symptom scores in individuals with lifetime MDD were significantly higher, whereas that in individuals without lifetime MDD were lower compared with the full study sample. Of note, a small subset of participants ($n = 2191$) completed the online mental health questionnaire up to 1.8 years after the neuroimaging session.

Inflammation Indices

Peripheral inflammation was indexed by baseline serum CRP level (mg/L), which was measured by an immunoturbidimetric high-sensitivity analysis on a Beckman Coulter AU5800 (see biomarker assay quality procedures for more details: https://biobank.ctsu.ox.ac.uk/showcase/showcase/docs/biomarker_issues.pdf). The blood sample used for CRP assessment was collected at the baseline visit, which was on average 7.6 years prior to the acquisition of DWI data and depression assessment in this study sample.

We also generated indices of neuroinflammation using DWI data (see details below in [DBSI and Neuroinflammation Index](#)).

Imaging Acquisition and Processing

This study made use of DWI and T1-weighted structural MRI data (i.e., FreeSurfer-based segmentation) that were processed and generated by an image-processing pipeline developed and run on behalf of UKB (50). Specifically, DWI data ($2 \times 2 \times 2 \text{ mm}^3$) were acquired using a multishell

approach with 2 b values ($b_1 = 1000 \text{ s/mm}^2$; $b_2 = 2000 \text{ s/mm}^2$). For each diffusion-weighted shell, 50 diffusion-encoding directions were acquired. Preprocessing of DWI data included eddy currents and head motion corrections, outlier slice correction, and gradient distortion correction (50).

High-resolution T1-weighted structural MRI data ($1 \times 1 \times 1 \text{ mm}^3$) was acquired with an in-plane acceleration sequence. The preprocessed T1-weighted images (i.e., after removing the skull and correcting for gradient distortion) were further processed with FreeSurfer, and the outputs (e.g., images, surface files, and summary outputs) were made available for download from the UKB Data Showcase. Details of the acquisition protocols, image processing pipeline, and derived imaging measures can be found in the UKB Imaging Documentation (<https://www.fmrib.ox.ac.uk/ukbiobank/>) and Miller *et al.* (51).

DBSI and Neuroinflammation Index

To index inflammation in the brain (neuroinflammation), we derived RF from DWI data using DBSI (36,39,41).

We focused on the hippocampus and amygdala, our regions of interest, and obtained subject-specific segmentation images of these 2 subcortical regions from the UKB FreeSurfer outputs, which were generated by FreeSurfer's *aseg* tool (52). We then extracted mean DBSI-RF from these regions of interest, separately for the left and right hemispheres, using the FreeSurfer segments as masks and FSL function *fsImeants* (53). The extracted RF values were then averaged within each ROI because they showed moderate-to-high left-right correlations ($r_s > 0.68$). Additionally, we derived a global index of neuroinflammation in whole-brain gray matter (i.e., global-RF), using the FSL FAST-based gray matter segmentation (54) as the mask. The resulting RFs of the whole-brain gray matter, hippocampus, and amygdala were included in separate statistical analyses.

Covariates

Age, sex, body mass index, ethnicity, smoking, and drinking statuses were included as covariates to adjust for potential

confounds. Additionally, we included a composite measure to index medical conditions that combined information about cancer diseases, noncancer diseases, medical operations, and medications. Covariate measurements and their variable IDs in the UKB database are described in [Table S1](#).

Statistical Analysis

To investigate whether DBSI-RF relates to depression symptoms, we conducted a series of linear regression analyses. Whenever significant associations were observed between DBSI-RF, depression, and CRP, we conducted planned mediation analyses to estimate whether DBSI-RF may indirectly link CRP to depression, using the PROCESS macro modeling tool that is implemented in R [i.e., R package *bruceR* (55)]. These analyses were performed separately for whole-brain gray matter, hippocampus, and amygdala, with adjustment for covariates (i.e., adjusted for both dependent and mediator variables). In case of significant effects of neuroinflammation indices, we further tested in the follow-up regression analyses to determine whether their effects were independent of CRP concentration. In mediation analyses, CRP concentration and RDSs were included as independent and dependent variables, respectively. We conducted these analyses for the full study sample, as well as for the selected subset of participants based on lifetime MDD definition. The regression models for lifetime MDD participants further included an interaction term of lifetime MDD status (“Yes” vs. “No”) with DBSI-RFs. To ensure robust statistical results, we conducted diagnostic analyses to test multicollinearity of independent variables in each regression model that returned significant effects of interest. Specifically, we calculated a generalized variance inflation factor score for each variable under investigation and investigated whether any observed effects might be biased due to high generalized variance inflation factor values (i.e., high collinearity).

False discovery rate (FDR) corrections were applied to all regression analyses to account for multiple testing.

RESULTS

Study Sample Demographics

Participants were predominately White and middle-old aged (mean age = 62.41 years), and about half were female. On average, participants scored 5.15 (SD = 1.68) on RDS questions, indicating relatively low levels of depression. Lifetime MDD participants showed similar demographic characteristics as the full study sample. Detailed demographic information for the study samples is summarized in [Table 1](#).

Associations Between Neuroinflammation, Peripheral Inflammation, and Depression

Peripheral Inflammation and Depression. Consistent with a recent study leveraging a larger UKB sample unconstrained by the need for neuroimaging data (56), CRP was positively associated with RDSs in our study sample ($t_{11499} = 3.45$, $\beta = 0.04$, false discovery rate-corrected p [p_{FDR}] < .005). This association remained following covariate inclusion ($t_{11490} = 2.57$, $\beta = 0.02$, $p_{FDR} < .05$).

Associations With Neuroinflammation Estimates.

DBSI-RF metrics in whole-brain gray matter, hippocampus, and amygdala were associated with each other ($\beta = 0.09 - 0.41$, $p_{FDRS} < .005$), with the strongest correlation between amygdala-RF and hippocampus-RF. Interestingly, only global-RF was associated with CRP concentration ($t_{11499} = -3.64$, $\beta = -0.03$, $p_{FDR} < .001$); this negative association remained significant with adjustment for all covariates ($t_{11490} = -2.08$, $\beta = -0.02$, $p_{FDR} < .05$). We did not find an association of hippocampus-RF ($\beta = -0.01$, $p_{FDR} = .17$), or of amygdala-RF ($\beta = -0.01$, $p_{FDR} = .12$) with CRP.

In line with our hypotheses, RF of the amygdala was associated with significantly more depressive symptoms with the adjustment for covariates ($t_{11490} = 2.53$, $\beta = 0.02$, $p_{FDR} < .05$) ([Figure 1A](#)) and further inclusion of CRP concentration ($t_{11489} = 2.58$, $\beta = 0.02$, $p_{FDR} < .05$). Neither the global-RF ($t_{11490} = 1.87$, $\beta = 0.02$, $p_{FDR} = .10$) nor the hippocampus-RF ($t_{11490} = 1.23$, $\beta = 0.01$, $p_{FDR} = .21$) showed such an association. Summary statistics for these associations between neuroinflammation estimates and depressive symptoms can be found in [Table S3](#).

As CRP was only significantly associated with global-RF, we conducted the hypothesized mediation model for these variables. However, we did not find evidence supporting a mediation role of global-RF on the association between peripheral inflammation and depression (indirect effect ab [here refers to the measure of the amount of mediation, with a indicating the path from variable X (i.e., peripheral inflammation) to variable M (i.e., neuroinflammation), and b the path from variable M to Y (i.e., depression)]: $Z = -1.33$, $p = .18$).

Secondary Analyses for Lifetime MDD Phenotype

Results from our secondary tests on lifetime MDD phenotype broadly recapitulated the findings observed with RDSs described above, with minor differences ([Table S4](#)).

Briefly, although we did not observe differences in CRP concentration or in global-RF or hippocampus-RF between lifetime MDD and non-lifetime MDD participants ($ps > .3$), we found a significant interactive effect of group (lifetime MDD vs. no lifetime MDD) and amygdala-RF on depression symptoms ($t_{6947} = 2.59$, $\beta = 0.08$, $p_{FDR} < .05$) ([Figure 1B](#)). This effect was robust against all covariates and further inclusion of CRP concentration ($t_{6946} = 2.6$, $\beta = 0.08$, $p_{FDR} < .05$). The follow-up tests showed that amygdala-RF had a stronger relation to depression symptoms in individuals with ($t_{1682} = 3.02$, $\beta = 0.07$, $p < .005$) than in those without lifetime MDD ($t_{5256} = 2.32$, $\beta = 0.03$, $p < .05$). These effects remained with further adjustment for CRP effect ($ps < .03$).

Robustness of Effects

Because some of the predictors and covariates were correlated ([Figure S1](#)), we conducted follow-up tests to examine whether the collinearity between predictors and covariates might have biased the observed results. Our findings show that the observed associative effects of CRP, and DBSI-RF of the amygdala on depression symptoms were robust against multicollinearity, as indicated by very low generalized variance inflation factor values for each independent variable in the regression models (i.e., all below 1.12; see full list in [Table S5](#)).




Figure 1. Robust association between the amygdala inflammation index and depression. **(A)** Restricted fraction—the index of neuroinflammation—in the amygdala was associated with depression: It predicted recent depressive symptom severity in the full study sample **(B)** and showed a stronger predictive effect on participants with lifetime major depressive disorder (MDD) than those without lifetime MDD. Note, all results remained significant with the removal of the data point in the no Lifetime MDD group that showed an abnormally large value (more than 3 SD above mean) of amygdala-restricted fraction (see [Robustness of Effects](#)).

Additionally, because nearly half of the participants in the full study sample reported no symptoms ($n = 5592$)—"Not at all" to all 4 symptom questions—we further conducted a logistic regression analysis to test whether the neuroinflammation indices of the amygdala could also predict group identities (i.e., control group = reporting zero symptoms). We replicated the amygdala-RF effect, showing that amygdala neuroinflammation could differentiate 2 groups of participants based on their depressive symptoms ($Z = 3.06$, $\beta = 0.06$, $p < .005$), with specificity of 0.62 and sensitivity of 0.56. This predictive effect remained when CRP concentration was considered in addition to the inclusion of all covariates ($Z = 3.09$, $\beta = 0.06$, $p < .005$).

Furthermore, to ensure that the effects of amygdala-RF were not driven by outliers ([Figure 1](#)), we repeated the analyses for the full study sample and lifetime MDD, after excluding the outliers above 3 standard deviations of the sample mean ($n = 91$ in the full study sample; $n = 59$ in the lifetime MDD sample). Results remained consistent with minor changes in coefficients (main effect in full study sample: $t_{11399} = 2.96$, $\beta = 0.03$, $p < .005$; interactive effect with lifetime MDD label: $t_{6896} = 2.26$, $\beta = 0.06$, $p < .05$).

Finally, we repeated the analysis with the amygdala-RF by lifetime MDD status interaction with further consideration of the time difference between these 2 measurements, which was available for $n = 8528$ participants. The result remained significant with adjustment for all covariates, including this time lag, as well as CRP ($t_{8515} = 2.89$, $\beta = 0.03$, $p < .005$). Furthermore, we repeated the same analysis for a subset of participants ($n = 4769$), whose assessment of lifetime MDD phenotype was performed prior to that of RDSs (i.e., ensuring lifetime MDD as a history in time). We observed the same interaction effect ($t_{2031} = 2.86$, $\beta = 0.14$, $p < .005$), which was driven by significant amygdala-RF impact on depression symptomatology in individuals with lifetime MDD ($n = 842$; $t_{830} = 2.97$, $\beta = 0.11$, $p < .005$). Amygdala-RF showed no association with depression in individuals without lifetime MDD ($t_{1190} = 1.11$, $\beta = 0.02$, $p = .27$). When we further restricted

analyses to a subsample of participants whose time lag between lifetime MDD and imaging assessments occurred within 0 to 6 months of one another ($n = 587$), the lifetime MDD by amygdala-RF interaction remained significant ($t_{572} = 2.92$, $\beta = 0.23$, $p < .005$), which was again driven by the lifetime MDD group ($n = 230$; $t_{217} = 2.49$, $\beta = 0.17$, $p < .05$). Participants without lifetime MDD ($n = 357$) showed no association between amygdala-RF and depression ($t_{344} = -.22$, $\beta = -0.01$, $p = .82$).

DISCUSSION

In the current study, we applied a novel noninvasive imaging technique to derive index measures of neuroinflammation in 2 a priori-selected subcortical regions—the hippocampus and amygdala—that have been consistently linked to depression symptomatology (29–31,33). Our study aimed to investigate whether the derived neuroinflammation indices could explain variance in depressive symptom severity and whether such neuroinflammation might mediate the impact of peripheral inflammation on depression. Our findings demonstrate an association between neuroinflammation at brain regional levels and depressive symptomatology.

In particular, our findings show that the putative neuroinflammation index of the amygdala has a small but significant association with depressive symptoms, especially in individuals with a lifetime history of MDD. This signal in the amygdala retained its association with depression even after adjusting for effects of demographics, lifestyles, and CRP concentration, which also showed significant associations with depressive symptoms. In light of recent investigations on the UKB dataset suggesting that the more strictly defined lifetime MDD phenotype [i.e., based on Composite International Diagnostic Interview-Short Form (47)] may have higher genomic and behavioral relevance specific to major depression (57,58), our results highlight a direct link between neuroinflammation index (i.e., inflammation-related cellularity) in the amygdala and elevated depressive symptoms. Our findings are

also consistent with a growing body of evidence showing that inflammation can contribute to depression via the stress-induced remodeling of amygdalar structure and function (59). Future studies may further investigate whether the depression-associated putative neuroinflammation is accompanied by other alterations in structure and functions in this subcortical region and in connected regions known to be involved in depression and whether stress-related experiences may modulate such associations.

Contrary to our hypotheses, however, we did not observe an association between hippocampus-RF and depression symptomatology even though the DBSI-RFs of the hippocampus and amygdala were moderately associated ($\beta = 0.41$), and both of these subcortical regions are considered highly relevant to depression (29–31,33). These results are broadly in line with recent studies that linked peripheral inflammation markers (e.g., CRP) to brain structure and function and suggested that specific brain circuitry might be more vulnerable to inflammation impact in individuals with depression (27,60,61). It awaits to be tested whether other brain regions, such as regions from the reward circuitry (e.g., ventral striatum, caudate nucleus, and putamen), may exhibit greater sensitivity to depression-related inflammation. Alternatively, previous studies have linked peripheral inflammation markers with depression symptom clusters (62) or subtypes (63), suggesting that inflammatory underpinnings of depression may be better characterized by symptom dimensions (64). In this study, we used a sum score of 4 items to indicate the overall severity of RDSs. The limited number of questions potentially restricted explorations on depression with a multidimensional approach. Future investigations may consider using depression assessment instruments that allow for quantifying symptom dimensions and examining the associated inflammatory profiles.

In line with the literature (6,9), we also found a positive association between the baseline CRP concentration and depressive symptom severity. This effect remained significant after controlling for physical illness and demographic characteristics including age, sex, body mass index, and smoking and drinking status. In contrast to our expectations, however, we did not find evidence suggesting that DBSI-RFs in whole-brain gray matter, hippocampus, or amygdala could mediate the CRP relationships to depression symptomatology. In fact, the amygdala-RF and baseline CRP seemed to explain complementary variance in depression symptoms in the study samples (i.e., showing significant effects separately in one multiple linear regression model). Notably, the collection of blood samples for baseline CRP concentration took place on average 7.6 years before DWI data acquisition. This large time difference might have blurred the relationship between the inflammation measures in the peripheral and central nervous systems. Nevertheless, when we repeated the models, including this time difference as an additional covariate, the relationships of baseline CRP concentration and neuroinflammation in the amygdala with depression symptomatology remained significant (amygdala: $t_{11487} = 2.68$, $\beta = 0.02$; CRP: $t_{11487} = 2.62$, $\beta = 0.02$; $p < .01$).

Limitations

The major limitation of this study concerned the time differences in data acquisition. First, the blood sample for assessing

CRP was collected on average 7.6 years prior to the acquisition of imaging data and assessment of depression symptoms. Although CRP is stable over time and prospectively associated with depression in older adults (13,65), this large time difference may have attenuated associations of both depression and neuroinflammation with CRP. Second, the collection of imaging data and the online questionnaire assessing lifetime MDD occurred at different times, resulting in varying time lags between these measurements at the individual level (i.e., ranging from 906 days before to 641 days after imaging acquisition). This variation may further muddy the associative effects in the current study. Nevertheless, our post hoc tests show that the interaction effect of amygdala-RF and lifetime MDD status remained with the additional inclusion of this time lag in the model (see details in [Robustness of Effects](#)). Additional time-related complications may also be induced by the time differences between depression onset and the acquisition of variables, and a history of depression itself may further bias our estimation. Yet, only a very small number of participants ($n = 176$) in the current study sample had an ICD-10 diagnosis of depression. When repeating analyses excluding these participants to minimize potential confounds associated with disorder onset, we observed the same associative effect of amygdala-RF with depression ($n = 11,304$; $t_{11292} = 2.62$, $\beta = 0.02$, $p < .01$) and the same interaction effect with lifetime MDD ($n = 4747$; $t_{2009} = 3.01$, $\beta = 0.14$, $p < .005$). It should also be noted that the neuroinflammation measure (i.e., DBSI-RF) in our study only indirectly assesses inflammation-related cellularity. Although it has not been tested in individuals with depression, DBSI-RF has been linked to increased cell nuclei numbers and activated microglia in several neuroinflammatory conditions (36,37,40,41) and associated with Alzheimer's disease and obesity (38,39,42), for which neuroinflammation is hypothesized to play a critical role (66,67). Our finding of the amygdala-RF in relation to depressive symptoms demonstrates a potential contribution of neuroinflammation to depression symptomatology. Nevertheless, due to the lack of evidence linking DBSI-RF with peripheral CRP in the current study (e.g., an insignificant association of amygdala-RF with CRP), our findings should be interpreted in the context that DBSI-RF is an indirect and putative index of neuroinflammation. Finally, our study only examined this putative neuroinflammation index in a limited number of brain regions and only in relation to depressive symptoms. It remains to be tested whether the observed amygdala-RF effect is specific to depression and what specific symptoms of depression may be most relevant. Given evidence linking peripheral inflammation and reward-related neural circuitry (60,61,68,69) as well as clinical presentations involving anhedonia and lethargy (68,70,71), this remains a promising avenue for future work.

Conclusions

Our study applied the novel DBSI technique to derive regional putative neuroinflammation indices for whole-brain gray matter, hippocampus, and amygdala and linked them with depression symptomatology. Although we failed to find evidence supportive of mediating effects for these neuroinflammation measures on the association between peripheral inflammation and depression, we did observe a robust and

significant effect of putative neuroinflammation in the amygdala on depressive symptom severity across individuals with and without a lifetime history of MDD, but more strongly in the former group. Future studies are warranted to expand the focus of brain regions outside the hippocampus and amygdala and to assess whether neuroinflammation in other depression-related regions (e.g., in reward circuitry) can also explain variance in depression symptomatology. With the very first evaluation of DBSI-based neuroinflammation in relation to depression symptomatology, our study provides early evidence suggesting the significance of regional neuroinflammation in depression symptom severity. These data add to a growing body of research linking inflammation with depression and demonstrating potential underlying mechanisms for depression pathogenesis.

ACKNOWLEDGMENTS AND DISCLOSURES

WZ is supported by McDonnell Center for Systems Neuroscience at Washington University in St. Louis; SAE received funding from Washington University in St. Louis Diabetes Research Center Pilot & Feasibility Award (Award No. P30 DK020579), Neuroimaging Laboratory Research Center Innovation Matching Funds, and the Mallinckrodt Institute of Radiology Pilot Award. RB received support from National Institutes of Health (Grant Nos. R01AG061162, R21AA027827, R01 DA054750, and U01DA055367). JDB is supported by the National Institutes of Health (Grant Nos. National Institute of Neurological Disorders and Stroke R34 NS118618; National Institute of Mental Health R01 MH128286) and the McDonnell Center for Systems Neuroscience.

The authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, Missouri (WZ, SAE, TH, JDB); Department of Psychiatry, Washington University School of Medicine, St. Louis, Missouri (JR, SAE, DMB, TH); Department of Obstetrics and Gynecology, Washington University School of Medicine, St. Louis, Missouri (YW); Department of Electrical and Systems Engineering, Washington University, St. Louis, Missouri (YW); Department of Psychological & Brain Sciences, Washington University, St. Louis, Missouri (DMB, TH, RB); and Department of Neurology, Washington University School of Medicine, St. Louis, Missouri (TH).

RB and JDB contributed equally to this work as joint senior authors.

Address correspondence to Wei Zhang, Ph.D., at wzzang@me.com, or Ryan Bogdan, Ph.D., at rbogdan@wustl.edu.

Received Dec 22, 2022; revised Apr 3, 2023; accepted Apr 29, 2023.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.bpsc.2023.04.011>.

REFERENCES

- Shorey S, Ng ED, Wong CHJ (2022): Global prevalence of depression and elevated depressive symptoms among adolescents: A systematic review and meta-analysis. *Br J Clin Psychol* 61:287–305.
- Hasin DS, Sarvet AL, Meyers JL, Saha TD, Ruan WJ, Stohl M, Grant BF (2018): Epidemiology of adult DSM-5 major depressive disorder and its specifiers in the United States. *JAMA Psychiatry* 75:336–346.
- Beurel E, Toups M, Nemeroff CB (2020): The bidirectional relationship of depression and inflammation: Double trouble. *Neuron* 107:234–256.
- Mac Giollabhui N, Ng TH, Ellman LM, Alloy LB (2021): The longitudinal associations of inflammatory biomarkers and depression revisited: Systematic review, meta-analysis, and meta-regression. *Mol Psychiatry* 26:3302–3314.
- Liu Y, Ho RC-M, Mak A (2012): Interleukin (IL)-6, tumour necrosis factor alpha (TNF- α) and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive disorder: A meta-analysis and meta-regression. *J Affect Disord* 139:230–239.
- Dowlati Y, Herrmann N, Swardfager W, Liu H, Sham L, Reim EK, Lanctôt KL (2010): A meta-analysis of cytokines in major depression. *Biol Psychiatry* 67:446–457.
- Hiles SA, Baker AL, de Malmanche T, Attia J (2012): A meta-analysis of differences in IL-6 and IL-10 between people with and without depression: Exploring the causes of heterogeneity. *Brain Behav Immun* 26:1180–1188.
- Howren MB, Lamkin DM, Suls J (2009): Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosom Med* 71:171–186.
- Osimo EF, Pillinger T, Rodriguez IM, Khandaker GM, Pariante CM, Howes OD (2020): Inflammatory markers in depression: A meta-analysis of mean differences and variability in 5,166 patients and 5,083 controls. *Brain Behav Immun* 87:901–909.
- Miller AH, Raison CL (2016): The role of inflammation in depression: From evolutionary imperative to modern treatment target. *Nat Rev Immunol* 16:22–34.
- Au B, Smith KJ, Gariépy G, Schmitz N (2015): The longitudinal associations between C-reactive protein and depressive symptoms: Evidence from the English Longitudinal Study of Ageing (ELSA). *Int J Geriatr Psychiatry* 30:976–984.
- Gimeno D, Kivimäki M, Brunner EJ, Elovainio M, De Vogli RD, Steptoe A, et al. (2009): Associations of C-reactive protein and interleukin-6 with cognitive symptoms of depression: 12-year follow-up of the Whitehall II study. *Psychol Med* 39:413–423.
- Bondy E, Norton SA, Voss M, Marks RB, Boudreaux MJ, Treadway MT, et al. (2021): Inflammation is associated with future depressive symptoms among older adults. *Brain Behav Immun Health* 13:100226.
- Miller AH, Maletic V, Raison CL (2009): Inflammation and its discontents: The role of cytokines in the pathophysiology of major depression. *Biol Psychiatry* 65:732–741.
- Kuhlman KR, Robles TF, Dooley LN, Boyle CC, Haydon MD, Bower JE (2018): Within-subject associations between inflammation and features of depression: Using the flu vaccine as a mild inflammatory stimulus. *Brain Behav Immun* 69:540–547.
- Glaser R, Robles TF, Sheridan J, Malarkey WB, Kiecolt-Glaser JK (2003): Mild depressive symptoms are associated with amplified and prolonged inflammatory responses after influenza virus vaccination in older adults. *Arch Gen Psychiatry* 60:1009–1014.
- Köhler O, Benros ME, Nordentoft M, Farkouh ME, Iyengar RL, Mors O, Krogh J (2014): Effect of anti-inflammatory treatment on depression, depressive symptoms, and adverse effects: A systematic review and meta-analysis of randomized clinical trials. *JAMA Psychiatry* 71:1381–1391.
- Raison CL, Rutherford RE, Woolwine BJ, Shuo C, Schettler P, Drake DF, et al. (2013): A randomized controlled trial of the tumor necrosis factor antagonist infliximab for treatment-resistant depression: The role of baseline inflammatory biomarkers. *JAMA Psychiatry* 70:31–41.
- Rapaport MH, Nierenberg AA, Schettler PJ, Kinkead B, Carudoos A, Walker R, Mischoulon D (2016): Inflammation as a predictive biomarker for response to omega-3 fatty acids in major depressive disorder: A proof-of-concept study. *Mol Psychiatry* 21:71–79.
- Copeland WE, Shanahan L, Worthman C, Angold A, Costello EJ (2012): Cumulative depression episodes predict later C-reactive protein levels: A prospective analysis. *Biol Psychiatry* 71:15–21.
- Troubat R, Barone P, Leman S, Desmidt T, Cressant A, Atanasova B, et al. (2021): Neuroinflammation and depression: A review. *Eur J Neurosci* 53:151–171.
- Hannestad J, DellaGioia N, Gallezot JD, Lim K, Nabulsi N, Esterlis I, et al. (2013): The neuroinflammation marker translocator protein is not elevated in individuals with mild-to-moderate depression: A [^{11}C]PBR28 PET study. *Brain Behav Immun* 33:131–138.
- Richards EM, Zanotti-Fregonara P, Fujita M, Newman L, Farmer C, Ballard ED, et al. (2018): PET radioligand binding to translocator protein (TSPO) is increased in unmedicated depressed subjects. *EJNMMI Res* 8:57.
- Holmes SE, Hinz R, Conen S, Gregory CJ, Matthews JC, Anton-Rodriguez JM, et al. (2018): Elevated translocator protein in anterior

- cingulate in major depression and a role for inflammation in suicidal thinking: A positron emission tomography study. *Biol Psychiatry* 83:61–69.
25. Yrondi A, Auouizerate B, El-Hage W, Moliere F, Thalamos C, Delcourt N, *et al.* (2018): Assessment of translocator protein density, as marker of neuroinflammation, in major depressive disorder: A pilot, multicenter, comparative, controlled, brain PET study (INFLADEP study). *Front Psychiatry* 9:326.
 26. Marek S, Tervo-Clemmens B, Calabro FJ, Montez DF, Kay BP, Hatoum AS, *et al.* (2022): Reproducible brain-wide association studies require thousands of individuals. *Nature* 603:654–660.
 27. Mehta ND, Haroon E, Xu X, Woolwine BJ, Li Z, Felger JC (2018): Inflammation negatively correlates with amygdala-ventromedial prefrontal functional connectivity in association with anxiety in patients with depression: Preliminary results. *Brain Behav Immun* 73:725–730.
 28. Setiawan E, Wilson AA, Mizrahi R, Rusjan PM, Miller L, Rajkowska G, *et al.* (2015): Role of translocator protein density, a marker of neuroinflammation, in the brain during major depressive episodes. *JAMA Psychiatry* 72:268–275.
 29. Schmaal L, Pozzi E, Ho CT, van Velzen LS, Veer IM, Opel N, *et al.* (2020): ENIGMA MDD: seven years of global neuroimaging studies of major depression through worldwide data sharing. *Transl Psychiatry* 10:1–19.
 30. Videbech P, Ravnkilde B (2004): Hippocampal volume and depression: A meta-analysis of MRI studies. *Am J Psychiatry* 161:1957–1966.
 31. Nolan M, Roman E, Nasa A, Levins KJ, O'Hanlon E, O'Keane V, William Roddy D (2020): Hippocampal and amygdala volume changes in major depressive disorder: A targeted review and focus on stress. *Chronic Stress (Thousand Oaks)* 4:2470547020944553.
 32. Spellman T, Liston C (2020): Toward circuit mechanisms of pathophysiology in depression. *Am J Psychiatry* 177:381–390.
 33. Park C, Rosenblat JD, Lee Y, Pan Z, Cao B, Iacobucci M, McIntyre RS (2019): The neural systems of emotion regulation and abnormalities in major depressive disorder. *Behav Brain Res* 367:181–188.
 34. Dillon DG, Pizzagalli DA (2018): Mechanisms of memory disruption in depression. *Trends Neurosci* 41:137–149.
 35. Hammen C (2018): Risk factors for depression: An autobiographical review. *Annu Rev Clin Psychol* 14:1–28.
 36. Wang Y, Wang Q, Haldar JP, Yeh FC, Xie M, Sun P, *et al.* (2011): Quantification of increased cellularity during inflammatory demyelination. *Brain* 134:3590–3601.
 37. Cross AH, Song SK (2017): A new imaging modality to non-invasively assess multiple sclerosis pathology. *J Neuroimmunol* 304:81–85.
 38. Samara A, Murphy T, Strain J, Rutlin J, Sun P, Neyman O, *et al.* (2019): Neuroinflammation and white matter alterations in obesity assessed by diffusion basis spectrum imaging. *Front Hum Neurosci* 13:464.
 39. Wang Q, Wang Y, Liu J, Sutphen CL, Cruchaga C, Blazey T, *et al.* (2019): Quantification of white matter cellularity and damage in preclinical and early symptomatic Alzheimer's disease. *NeuroImage Clin* 22:101767.
 40. Strain JF, Burdo TH, Song SK, Sun P, El-Ghazzawy O, Nelson B, *et al.* (2017): Diffusion basis spectral imaging detects ongoing brain inflammation in virologically well-controlled HIV+ patients. *J Acquir Immune Defic Syndr* 76:423–430.
 41. Wang Y, Sun P, Wang Q, Trinkaus K, Schmidt RE, Naismith RT, *et al.* (2015): Differentiation and quantification of inflammation, demyelination and axon injury or loss in multiple sclerosis. *Brain* 138:1223–1238.
 42. Ly M, Raji CA, Yu GZ, Wang Q, Wang Y, Schindler SE, *et al.* (2021): Obesity and white matter neuroinflammation related edema in Alzheimer's disease dementia biomarker negative cognitively normal individuals. *J Alzheimers Dis* 79:1801–1811.
 43. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, *et al.* (2015): UK Biobank: An open access resource for identifying the causes of a Wide Range of complex diseases of middle and old age. *PLoS Med* 12:e1001779.
 44. Ansar W, Ghosh S (2013): C-reactive protein and the biology of disease. *Immunol Res* 56:131–142.
 45. Dutt RK, Hannon K, Easley TO, Griffis JC, Zhang W, Bijsterbosch JD (2022): Mental health in the UK Biobank: A roadmap to self-report measures and neuroimaging correlates. *Hum Brain Mapp* 43:816–832.
 46. Davis KAS, Coleman JRI, Adams M, Allen N, Breen G, Cullen B, *et al.* (2020): Mental health in UK Biobank – Development, implementation and results from an online questionnaire completed by 157 366 participants: A reanalysis. *BJPsych Open* 6:e18.
 47. Kessler RC, Andrews G, Mroczek D, Ustun B, Wittchen H-U (1998): The World Health Organization Composite International Diagnostic Interview short-form (CIDI-SF). *Int J Method Psychiat Res* 7:171–185.
 48. Howard DM, Adams MJ, Shirali M, Clarke TK, Marioni RE, Davies G, *et al.* (2018): Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways. *Nat Commun* 9:1470.
 49. Coleman JRI, Peyrot WJ, Purves KL, Davis KAS, Rayner C, Choi SW, *et al.* (2020): Genome-wide gene-environment analyses of major depressive disorder and reported lifetime traumatic experiences in UK Biobank. *Mol Psychiatry* 25:1430–1446.
 50. Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Griffanti L, Douaud G, *et al.* (2018): Image processing and quality control for the first 10,000 brain imaging datasets from UK Biobank. *NeuroImage* 166:400–424.
 51. Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, *et al.* (2016): Multimodal population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci* 19:1523–1536.
 52. Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, *et al.* (2002): Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron* 33:341–355.
 53. Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM (2012). *FSL*. *NeuroImage* 62:782–790.
 54. Zhang Y, Brady M, Smith S (2001): Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* 20:45–57.
 55. Bao H-W-S (2022): *bruceR: Broadly useful convenient and efficient R functions*. Available at: <https://CRAN.R-project.org/package=bruceR>. Accessed October 10, 2022.
 56. Pitharoulis MC, Hagenars SP, Glanville KP, Coleman JRI, Hotopf M, Lewis CM, Pariante CM (2021): Elevated C-reactive protein in patients with depression, independent of genetic, health, and psychosocial factors: Results from the UK Biobank. *Am J Psychiatry* 178:522–529.
 57. Cai N, Revez JA, Adams MJ, Andlauer TFM, Breen G, Byrne EM, *et al.* (2020): Minimal phenotyping yields genome-wide association signals of low specificity for major depression. *Nat Genet* 52:437–447.
 58. de Nooij L, Harris MA, Adams MJ, Clarke TK, Shen X, Cox SR, *et al.* (2020): Cognitive functioning and lifetime major depressive disorder in UK Biobank. *Eur Psychiatry* 63:e28.
 59. Hu P, Lu Y, Pan BX, Zhang WH (2022): New insights into the pivotal role of the amygdala in inflammation-related depression and anxiety disorder. *Int J Mol Sci* 23:11076.
 60. Felger JC, Haroon E, Patel TA, Goldsmith DR, Wommack EC, Woolwine BJ, *et al.* (2020): What does plasma CRP tell us about peripheral and central inflammation in depression? *Mol Psychiatry* 25:1301–1311.
 61. Felger JC (2018): Imaging the role of inflammation in mood and anxiety-related disorders. *Curr Neuropharmacol* 16:533–558.
 62. Franklyn SI, Stewart J, Beaurepaire C, Thaw E, McQuaid RJ (2022): Developing symptom clusters: Linking inflammatory biomarkers to depressive symptom profiles. *Transl Psychiatry* 12:133.
 63. Moriarity DP, Slavich GM, Alloy LB, Olino TM (2023): Hierarchical inflammatory phenotypes of depression: A novel approach across five independent samples and 27,730 adults. *Biol Psychiatry* 93:253–259.
 64. Bekhbat M, Li Z, Mehta ND, Treadway MT, Lucido MJ, Woolwine BJ, *et al.* (2022): Functional connectivity in reward circuitry and symptoms of anhedonia as therapeutic targets in depression with high inflammation: Evidence from a dopamine challenge study. *Mol Psychiatry* 27:4113–4121.

Amygdala Neuroinflammation Associated With Depression

65. Sonsin-Diaz N, Gottesman RF, Fracica E, Walston J, Windham BG, Knopman DS, Walker KA (2020): Chronic systemic inflammation is associated with symptoms of late-life depression: The ARIC study. *Am J Geriatr Psychiatry* 28:87–98.
66. Leng F, Edison P (2021): Neuroinflammation and microglial activation in Alzheimer disease: Where do we go from here? *Nat Rev Neurol* 17:157–172.
67. Miller AA, Spencer SJ (2014): Obesity and neuroinflammation: A pathway to cognitive impairment. *Brain Behav Immun* 42:10–21.
68. Costi S, Morris LS, Collins A, Fernandez NF, Patel M, Xie H, *et al.* (2021): Peripheral immune cell reactivity and neural response to reward in patients with depression and anhedonia. *Transl Psychiatry* 11:565.
69. Kudinova AY, Gano A, James KM, Lawlor C, Deak T, Gibb BE (2020): Anhedonia and increased evoked immune response. *Brain Behav Immun Health* 6:100090.
70. Felger JC, Treadway MT (2017): Inflammation effects on motivation and motor activity: Role of dopamine. *Neuropsychopharmacology* 42:216–241.
71. Eisenberger NI, Berkman ET, Inagaki TK, Rameson LT, Mashal NM, Irwin MR (2010): Inflammation-induced anhedonia: Endotoxin reduces ventral striatum responses to reward. *Biol Psychiatry* 68:748–754.