

**Table 21. Weed Response to Herbicides in Grass Pastures/CRP/Grass Hay**

This table compares the relative effectiveness of herbicides on weeds. Ratings are based on labeled application rate and weed size or growth stage. Control of perennial weeds may require more than one application. Performance may be better or worse than indicated in the table, due to weather or soil conditions, or other variables.

Weed control rating:  
 9 = 90% to 100% control  
 8 = 80% to 90% control  
 7 = 70% to 80% control

insufficient information  
 6 = 60% to 70% control  
 - = not recommended or

	2,4-D	Chaparral	Cimarron Max	Clopyralid	Crossbow	Curtail	Dicamba	GrazonNext HL	Glyphosate	Metsulfuron	Milestone	PastureGard	Remedy Ultra
Site of action	4	2/4	2/4	4	4	4	4	4	9	2	4	4	4
<b>Winter annual</b>													
Chamomile, mayweed	7	8	9	9	7	9	8	7	9	9	-	-	-
Chickweed, common	-	9	9	6	8	6	9	8	9	9	7	8	8
Cockle, corn	7	-	9	-	8	7	9	6	9	-	-	-	-
Cockle, cow	7	-	9	-	8	7	9	6	9	-	-	-	-
Horseweed (marestail)	9	9	9	9	9	9	9	9	9	9	8	6	7
Mustard spp.	9	8	9	6	9	9	7	9	9	9	-	-	9
Pennycress, field	9	8	9	6	9	9	8	9	9	9	-	-	-
Pepper weed spp.	9	8	9	6	9	9	7	9	9	9	-	-	-
Shepherd's-purse	8	8	9	6	9	8	7	8	9	9	-	-	-
<b>Summer annual</b>													
Cocklebur, common	8	9	9	9	9	9	9	9	9	9	9	8	8
Lambsquarters, common	8	9	9	6	9	8	9	9	9	9	9	9	9
Nightshade, black	7	9	8	-	8	7	8	9	8	7	9	-	-
Pigweed spp.	8	9	9	6	8	8	9	8	9	9	8	9	-
Ragweed, common	9	9	9	9	9	9	9	9	9	-	9	9	9
Ragweed, giant	9	9	9	9	9	9	9	9	9	-	9	9	9
Velvetleaf	8	7	9	-	8	8	9	8	8	8	-	9	-
<b>Biennial</b>													
Burdock, common	9	9	9	8	9	8	7	9	9	9	9	9	9
Carrot, wild	8	9	9	6	9	8	7	8	8	9	-	7	8
Evening primrose, common	8	8	8	-	9	7	7	9	9	-	8	-	8
Hemlock, poison	7	-	8	-	9	8	8	7	8	-	-	-	9
Lettuce, wild	9	9	9	9	8	9	8	9	8	9	-	-	-
Parsnip, wild	8	9	8	-	9	8	8	8	8	8	-	-	-
Teasel	7	9	9	8	8	9	8	8	8	9	-	-	-
Thistle, bull	8	9	9	9	9	9	9	9	9	9	9	6	6
Thistle, musk	9	9	9	9	9	9	9	9	9	9	9	6	7
Yellow rocket	8	8	9	-	9	8	8	8	8	9	-	-	-