

safe-guard[®] AquaSol[®] (fenbendazole oral suspension) for Poultry

Innovative manufacturing process enables easy, effective deworming for broilers, breeders and pullets

NEW PRODUCTION TECHNOLOGY

- Proprietary, innovative wet-milling technique produces **highly stable suspension** in drinking water
- Smaller particle size helps **prevent sedimentation** and clogged pipes, filters or nipple drinkers
- Easily administered by **medicators or proportioners**

APPROVED FOR MAJOR INTESTINAL PARASITES

- *Ascaridia galli* (adult stage roundworms)
- *Heterakis gallinarum* (adult stage cecal worms)

EFFECTIVE

- **98.8% efficacy** against *A. galli* in 4 US field studies (**100% in broilers**)
- **92.1% efficacy** against *H. gallinarum* (long lived birds)

CONVENIENT WATER FORMULATION

- Short, **5-day treatment**
- **No pre-slaughter withdrawal period** when used as directed
- **Shelf life of 3 years unopened, 6 months after opening** — multi-use container reduces waste

SAFE, PALATABLE

- **Proven safe** when used as directed
- **No adverse effects** on egg production, hatchability or progeny of breeders
- **No impact on water or feed intake** compared to unmedicated controls



Safe-Guard[®] AquaSol is available in 1 liter and 1 gallon presentations.

Product Code:
1 liter 148720 | 1 gallon 137151



safe-guard[®] AquaSol

(fenbendazole oral suspension) for Poultry

Suspension Concentrate, Antiparasitic

200 mg of fenbendazole/mL

For oral administration via drinking water

For Veterinary Use Only.

Description

Safe-Guard[®] AquaSol is a suspension concentrate containing fenbendazole, an antiparasitic. Each mL of Safe-Guard AquaSol contains 200 mg of fenbendazole.

Indications for Use

Safe-Guard AquaSol is indicated for the treatment and control of adult *Ascaridia galli* in broiler chickens and replacement chickens intended to become breeding chickens and for the treatment and control of adult *A. galli* and *Heterakis gallinarum* in breeding chickens.

Not for use in laying hens and replacement chickens intended to become laying hens.

Dosage and Administration

Safe-Guard AquaSol must be administered orally to chickens via the drinking water at a daily dose of 1.0 mg/kg BW (0.454 mg/lb) for 5 consecutive days.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Residue Warning

No withdrawal period is required when used according to labeling. Do not use the drug in laying hens.

Storage Information

Store at room temperature 30°C (86°F). Once opened, do not store the container above 25°C (77°F). Do not freeze. Protect from light.

Use within 6 months after opening. Use the medicated water within 24 hours.

For customer service, adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573