

The Asymptotic Weak Gravity Conjecture in M-theory

- w/ Cesar Cota, Alessandro Mininno, Max Wiesner
2212.09758 and 2208.00009

Timo Weigand

Cluster of Excellence Quantum Universe, Hamburg University

Motivation

Swampland Conjectures: profound predictions for *Strings and Geometry*

General conjectures on quantum gravity tested in string compactifications

This talk:

Tower Weak Gravity Conjecture

Specifically:

- Goal:

Show Tower WGC in M-theory on arbitrary CY3 for gauge groups with a weak coupling limit

⇒ **Asymptotic Tower WGC**

- Mathematical Input:

1. Kähler geometry of CY3
2. Counting of BPS *and of non-BPS states* via DT-invariants on CY3

Tower Weak Gravity Conjecture

initiated in [Arkani-Hamed,Motl,Nicolis,Vafa'06]

*Consider a gauge theory coupled to quantum gravity, (for simplicity) with abelian gauge factors $U(1)$ and charge lattice $\Lambda_{\mathbf{Q}}$. Then every ray in the lattice $\Lambda_{\mathbf{Q}}$ must support a **tower** of **super-extremal** states.*

Two notions of **super-extremality**:

1. with respect to extremal black hole:

$$\frac{g_{\text{YM}}^2 q^2}{m^2} \geq \frac{g_{\text{YM}}^2 Q^2}{M^2} \Big|_{\text{B.H.}}$$

2. self-repulsive:

$$F_{\text{Coulomb}} \geq F_{\text{Grav.}} + F_{\text{Yukawa}}$$

F_{Yukawa} in presence of massless scalars - first pointed out in [Palti,'17]

In general both notions are not equivalent [Heidenreich,Reece,Rudelius,'19]

but in asymptotic weak coupling limit they are [Lee,Lerche,TW'18]

Tower Weak Gravity Conjecture

initiated in [Arkani-Hamed,Motl,Nicolis,Vafa'06]

Consider a gauge theory coupled to quantum gravity, (for simplicity) with abelian gauge factors $U(1)$ and charge lattice $\Lambda_{\mathbf{Q}}$. Then every ray in the lattice $\Lambda_{\mathbf{Q}}$ must support a *tower* of *super-extremal* states.

Two notions of **super-extremality**:

1. with respect to extremal black hole:

$$\frac{g_{\text{YM}}^2 q^2}{m^2} \geq \frac{g_{\text{YM}}^2 Q^2}{M^2} \Big|_{\text{B.H.}}$$

2. self-repulsive:

$$F_{\text{Coulomb}} \geq F_{\text{Grav.}} + F_{\text{Yukawa}}$$

Tower: [Heidenreich,Reece,Rudelius'15-16] [Montero,Shiu,Soler'16] [Andriolo et al.'16]

\exists super-extremal particle of charge $n\mathbf{Q}$ for $n \in \mathcal{I}$ an *infinite* set

WGC: BPS Towers

Fact: Every BPS state is automatically super-extremal

⇒ Existence of tower of BPS states along ray in charge lattice sufficient for tower WGC in that direction

[Grimm,Palti,Valenzuela'18] [Gendler,Valenzuela'20] [Grimm,Heisteeg'20] . . .

[Alim,Heidenreich,Rudelius'21] [Gendler,Heidenreich,McAllister,Moritz,Rudelius'22]

Challenge for tower WGC:

What if there is no BPS tower in a certain direction in charge lattice?

Examples:

F-theory compactification

- to 6d $N = (1, 0)$ [Lee,Lerche,TW'18]
- to 4d $N = 1$ [Lee,Lerche,TW'19] [Kläwer, Lee, TW, Wiesner'20]
[Heidenreich,Reece,Rudelius'21] [Cota,Mininno,TW,Wiesner'22 (1)]

M-theory compactification to 5d $N = 1$ [Alim,Heidenreich,Rudelius'21]

[Cota,Mininno,TW,Wiesner'22 (2)] [Gendler,Heidenreich,McAllister,Moritz,Rudelius'22]

Beyond BPS Towers

Main result: [Cota, Mininno, TW, Wiesner'22 (2)]

Consider M-theory compactified on CY_3 to 5d $N = 1$.

Whenever a direction in the charge lattice does not admit a (super-extremal) BPS tower, then

- either there is no weak coupling limit for the relevant $U(1)$ s
- or there does exist a super-extremal non-BPS tower.

\implies Establishes *asymptotic* tower WGC in 5d M-theory

WGC in 5d M-theory

M-theory on CY3 X_3 : 5d N=1 theory (8 supercharges)

- Basis of **U(1) gauge groups** from expanding

$$C_3 = A^\alpha \wedge J_\alpha, \quad \alpha = 1, \dots, h^{1,1}(X_3)$$

- J_α : Basis of Kähler cone generators $J = v^\alpha J_\alpha$
- Gauge kinetic terms

$$S_{5d} = \frac{M_{\text{Pl}}^3}{2} \int_{\mathbb{R}^{1,4}} R \star 1 - \frac{1}{2g_5^2} \int_{\mathbb{R}^{1,4}} f_{\alpha\beta} F^\alpha \wedge \star F^\beta + \dots \quad F^\alpha = dA^\alpha$$

Gauge kinetic matrix $f_{\alpha\beta} \iff$ Kähler moduli (modulo overall volume)

$$f_{\alpha\beta} = \frac{1}{\mathcal{V}^{1/3}} \int_{X_3} J_\alpha \wedge \star J_\beta = \left(\hat{\mathcal{V}}_\alpha \hat{\mathcal{V}}_\beta - \hat{\mathcal{V}}_{\alpha\beta} \right)$$

$$\mathcal{V} = \frac{1}{6} \int_{X_3} J^3, \quad \mathcal{V}_\alpha = \frac{1}{2\mathcal{V}} \int_{X_3} J_\alpha \wedge J^2, \quad \mathcal{V}_{\alpha\beta} = \frac{1}{\mathcal{V}} \int_{X_3} J_\alpha \wedge J_\beta \wedge J, \quad \hat{v}^\alpha = \frac{v^\alpha}{\mathcal{V}}$$

WGC in 5d M-theory

Self-repulsiveness condition for states of

- charges Q_α under $U(1)_\alpha$
- Kähler moduli dependent mass $M_k(v^\alpha)$

$$\begin{aligned}
 F_{\text{Coulomb}} &\stackrel{!}{\geq} F_{\text{grav}} + F_{\text{Yukawa}} \\
 \frac{(M_{\text{Pl}} g_5^2)(Q_\alpha f^{\alpha\beta} Q_\beta)}{M_k^2/M_{\text{Pl}}^2} &\stackrel{!}{\geq} \frac{d-3}{d-2} \Big|_{d=5} + \frac{1}{2} \frac{M_{\text{Pl}}^4}{M_k^4} \left(f^{\alpha\beta} - \frac{1}{3} \hat{v}^\alpha \hat{v}^\beta \right) \partial_\alpha \left(\frac{M_k^2}{M_{\text{Pl}}^2} \right) \partial_\beta \left(\frac{M_k^2}{M_{\text{Pl}}^2} \right)
 \end{aligned}$$

basis of U(1) gauge groups from expanding

$$C_3 = A^\alpha \wedge J_\alpha, \quad \alpha = 1, \dots, h^{1,1}(X_3)$$

J_α : Basis of Kähler cone generators $J = v^\alpha J_\alpha$

$$S_{5d} = \frac{M_{\text{Pl}}^3}{2} \int_{\mathbb{R}^{1,4}} R \star 1 - \frac{1}{2g_5^2} \int_{\mathbb{R}^{1,4}} f_{\alpha\beta} F^\alpha \wedge \star F^\beta + \dots$$

WGC: BPS Towers

First source for super-extremal tower:

BPS particles in 5d from M2-branes on holomorphic curves on X_3

Conjecture: [Alim,Heidenreich,Rudelius'21]

Every curve class $C \in \text{Mov}_1(X_3)$ supports a tower of BPS states, i.e.

$$N_{g=0}(nC) \neq 0 \quad \forall C \in \text{Mov}_1(X_3)$$

Recall: Movable curve cone $\text{Mov}_1(X_3)$ is dual to cone of effective divisors $\text{Eff}^1(X_3)$

✓ confirmed in many examples in [Alim,Heidenreich,Rudelius'21][Gendler et al.'22]

✓ Black hole extremality condition = BPS condition

for such $C \in \text{Mov}_1(X_3)$ [Alim,Heidenreich,Rudelius'21]

Challenge for tower WGC: What if there are no BPS towers?

Example: Conifold

Beyond BPS Towers

Main result: [Cota, Mininno, TW, Wiesner'22]

Whenever there is no BPS tower, then

- either there is no weak coupling limit for the U(1)s
- or there does exist a super-extremal non-BPS tower.

Strategy:

1. Characterise all **weak coupling limits** \iff Kähler geometry
2. Identify **towers of super-extremal BPS or non-BPS states** for U(1)s with a weak coupling limit
 \iff DT invariants/Noether-Lefschetz theory

Weak Coupling Limits

$$S_{5d} = \frac{M_{\text{Pl}}^3}{2} \int_{\mathbb{R}^{1,4}} R \star 1 - \frac{1}{2g_5^2} \int_{\mathbb{R}^{1,4}} f_{\alpha\beta} F^\alpha \wedge \star F^\beta + \dots$$

Gauge kinetic matrix $f_{\alpha\beta} \iff$ Kähler moduli (at fixed overall volume)

$$f_{\alpha\beta} = \frac{1}{\mathcal{V}^{1/3}} \int_{X_3} J_\alpha \wedge \star J_\beta = (\hat{\mathcal{V}}_\alpha \hat{\mathcal{V}}_\beta - \hat{\mathcal{V}}_{\alpha\beta})$$

1) Necessary condition for weak coupling limit:

Entries of $f_{\alpha\beta} \rightarrow \infty \iff$ Infinite distance limits (in Kähler moduli space at fixed overall volume \mathcal{V}) cf. [Heidenreich, Rudelius'20]

2) More precise criterion:

$$U(1)_C = c_\alpha U(1)^\alpha \quad \text{basis } U(1)^\alpha \iff A^\alpha \text{ in } C_3 = A^\alpha J_\alpha$$


$$\Lambda_{\text{WGC}}^2 (U(1)_C) = g_{\text{YM,C}}^2 M_{\text{Pl}}^3 = g_5^2 (c_\alpha f^{\alpha\beta} c_\beta) M_{\text{Pl}}^3$$

$$\frac{\Lambda_{\text{WGC}}^2 (U(1)_C)}{\Lambda_{\text{QG}}^2} \rightarrow 0 \quad \Lambda_{\text{QG}} = \Lambda_{\text{sp.}} : \text{species scale for limit}$$


Weak Coupling Limits

Characterisation of weak coupling limits: [Cota, Mininno, TW, Wiesner'22]

follows from classification of infinite distance limits in classical Kähler moduli space of X_3 at fixed volume [Lee, Lerche, TW'19]:



KK tower from M2 on T^2
 Effective decomp. $5d \rightarrow 6d$
 $\Lambda_{\text{sp.}} = M_{\text{Pl.}, 6d}$





heterotic/Type II string from M5 on fiber
 5d emergent heterotic/Type II string limit
 $\Lambda_{\text{sp.}} = \Lambda_{\text{sp.}, \text{string}}$

Weak Coupling Limits

Characterisation of weak coupling limits: [Cota, Mininno, TW, Wiesner'22]

In M-theory compactified on a Calabi–Yau X_3 , the only $U(1)$ s which admit a weak coupling limit are obtained as $U(1)_C = c_\alpha U(1)^\alpha$ for $C = c_\alpha \omega^\alpha \in H_2(X_3)$ a **curve class** with

1. $C = T^2$ - a generic **torus fiber** of X_3
2. $C \subset S$ for S a generic **K3 or T^4 fiber** of X_3 or a degenerate such fiber occuring at finite distance in the fiber moduli space.



Elliptic tower counting

Suppose X_3 admits fibration $\pi : T^2 \rightarrow B_2$ with **generic fiber** $T^2 \equiv \mathcal{E}$.

Unless X_3 also admits a K3 or T^4 surface:

only U(1) with a weak coupling limit: $U(1)_{\mathcal{E}} = c_{\alpha} U(1)^{\alpha}$ $\mathcal{E} = c_{\alpha} \omega^{\alpha}$

- Intuition:


' $g_{\text{YM},C}^2 M_{\text{Pl}} \sim \mathcal{V}_C^{-2}$ ' \mathcal{V}_C : 'dual' divisor to C

- $\mathcal{V}_{T^2} \sim \frac{1}{\lambda}$, $\mathcal{V}_{B_2} \sim \lambda \implies g_{\text{YM},\mathcal{E}}^2 M_{\text{Pl}} \sim \frac{1}{\lambda^2}$

- $\frac{\Lambda_{\text{sp, KK}}^3}{M_{\text{Pl}}^3} = \left(\frac{M_{\text{KK}}^3}{M_{\text{Pl}}^3} \right)^{\frac{n}{3+n}} \Big|_{n=1}$

cf. [Montero, Vafa, Valenzuela '22]

- $\frac{M_{\text{KK}}^3}{M_{\text{Pl}}^3} \sim \mathcal{V}_{T^2}^3 \sim \frac{1}{\lambda^3} \implies \frac{\Lambda_{\text{sp, KK}}^3}{M_{\text{Pl}}^3} \sim \frac{1}{\lambda^{3/4}}$




$$\frac{\Lambda_{\text{WGC}}^2 (U(1)_{\mathcal{E}})}{\Lambda_{\text{QG}}^2} \sim \frac{g_{\text{YM},\mathcal{E}}^2 M_{\text{Pl}}^3}{\Lambda_{\text{sp, KK}}^2} \sim \frac{1/\lambda^2}{1/\lambda^{1/2}} \sim \frac{1}{\lambda^{3/2}} \rightarrow 0 \quad \checkmark$$

Elliptic tower counting

Unless X_3 also admits a K3 or T^4 surface:

- *only* $U(1)_\mathcal{E}$ has a weak coupling limit
- All other fibral curves not weakly coupled:

$$\frac{g_{\text{YM}, C^f}^2 M_{\text{Pl}}^3}{\Lambda_{\text{sp, KK}}^2} \sim \frac{1/\lambda^{1/2}}{1/\lambda^{1/2}} \sim 1$$



Super-extremal towers:

- \exists **tower of BPS states** charged under $U(1)_\mathcal{E}$:
M2-branes wrapped n -times on T^2 fiber

$$N_{n\mathcal{E}}^0 = -\chi(X_3)$$

\implies super-extremal BPS tower \iff asymptotic tower WGC

Interpretation: KK tower for decompactification 5d to 6d

- **No BPS towers along other fibral curves**, and no non-BPS tower known

Elliptic tower counting

Super-extremal towers:

- \exists tower of BPS states charged under $U(1)_{\mathcal{E}}$:
M2-branes wrapped n -times on T^2 fiber

$$N_{n\mathcal{E}}^0 = -\chi(X_3)$$

\implies super-extremal BPS tower \iff asymptotic tower WGC

Interpretation: KK tower for decompactification 5d to 6d

- No BPS towers along other fibral curves, and no non-BPS tower known
 \iff corresponding gauge group does not become weakly coupled in asymptotic 6d theory (by assumption B_2 not rationally fibered)

cf. [Cota, Mininno, TW, Wiesner'22 (1)]

- Gauge groups 'from base' become 2-forms in asymptotic 6d theory

K3 tower counting

Suppose X_3 admits fibration $\rho : K3 \rightarrow \mathbb{P}^1$

The *only** $U(1)$ which can undergo a **weak coupling limit** is


$$U(1)_C = c_\alpha U(1)^\alpha \quad C = c_\alpha \omega^\alpha$$

for C a **curve in a generic K3 fiber** or in a special K3 fiber at finite distance in moduli space.

- $g_{\text{YM},C}^2 M_{\text{Pl}} \sim \frac{1}{\lambda^2}$
- Species scale set by tension of emergent heterotic string cf. [Dvali,Lüst'09]

[Marchesano,Melotti'22] [Castellano,Herraez,Ibanez'22]

$$\Lambda_{\text{sp}}^2 \sim M_{\text{het}}^2 \log \left(\frac{M_{\text{Pl}}}{M_{\text{het}}} \right), \quad M_{\text{het}}^2 \sim \frac{1}{\lambda^2} M_{11\text{d}}^2$$



$$\frac{\Lambda_{\text{WGC}}^2 (U(1))}{\Lambda_{\text{sp}}^2} \sim \frac{1/\lambda^2}{1/\lambda^2(1 + \log(\lambda))} \rightarrow 0$$


K3 tower counting

Weakly coupled: $U(1)_C$, $C \subset$ generic/finite distance deg. K3

Such curves lie in a lattice

$$\Lambda_{\mathbb{R}}^* = \Lambda_+^* \oplus \Lambda_-^*$$

$\Lambda_{\mathbb{R}}^*$ lattice of charges with respect to such U(1) of signature $(1, r)$, $r \leq 19$



- $C^2 \geq 0$: BPS tower exists

✓ such curves are movable inside K3 fiber and hence in movable cone

✓ in agreement with BPS index counting via modular forms

Gopakumar-Vafa invariants for M2-brane on $C \subset K3$: [Harvey, Moore'99], ...


$$N_C^{g=0} = c\left(\frac{C^2}{2}\right) \quad f(q) = \sum_{n=-1}^{\infty} c(n)q^n \quad \text{modular form}$$

Infinite tower on $nC \leftrightarrow C \cdot C \geq 0$ = non-contractible curves inside K3

K3 tower counting

Weakly coupled: $U(1)_C$, $C \subset$ generic/finite distance deg. K3

- $C^2 \geq 0$: BPS tower exists
- $C^2 < 0$: No BPS tower exists
✓ curves are rigid inside K3 fiber and hence not in movable cone



Claim: Tower of non-BPS states takes over [Cota, Mininno, TW, Wiesner'22 (2)]

- Special set of states from excitations of MSW-type heterotic string obtained by wrapping M5-brane on K3 fiber is super-extremal.
- Existence established via relation of elliptic genus and 4d BPS invariants in Type IIA on $CY3^a$

[Bouchard, Creutzig, Diaconescu, Doran, Quigley, Sheshmani16] [Pandharipande, Thomas'16]

^aAnalysis most explicitly in absence of multi-components fibers, but expect results to carry over more generally.

K3 tower counting

M-theory on $X_3 \times S_M^1$

bound state of M5-brane on K3

(winding, KK) number (r, n)

M2 brane on curve \mathbf{Q} on K3

Type IIA string on X_3

D4-D2-D0 bound state

of charge vector

$$\gamma = (r\Sigma_{\text{K3}}, \mathbf{Q}, n)$$

Can view these states as winding modes of heterotic string from M5 on K3 at KK level n and charge vector \mathbf{Q}

Special case $r = 1$:

By level matching can identify $n \leftrightarrow n_L = (\text{left-moving})$ excitation level of single heterotic string

4d BPS states of charge
vector $\gamma = (\Sigma_{\text{K3}}, \mathbf{Q}, n)$

\implies

existence of non-BPS string
excitations in 5d at level
 $n_L = n$ and charge \mathbf{Q}

Goal: Show existence of super-extremal state at $n = -\frac{1}{2}\mathbf{Q}^2$

K3 tower counting

Modified elliptic genus of wrapped heterotic string [Gaiotto, Strominger, Yin'06]:

$$\begin{aligned} Z_{\mathbf{S}}^{(r=1)}(\tau, \bar{\tau}, \mathbf{z}) &= \text{Tr}_{\text{RR}} F_R^2 (-1)^{F_R} q^{L_0 - \frac{c_L}{24}} \bar{q}^{\bar{L}_0 - \frac{c_R}{24}} e^{2\pi i z^i Q_i} \\ &= \sum_{n^{(L)}, n^{(R)}} N(n^{(L)}, n^{(R)}, \mathbf{Q}) q^{n^{(L)} - 1} \bar{q}^{n^{(R)}} e^{2\pi i z^i Q_i} \end{aligned}$$

Expression in terms of 4d BPS numbers:

$$Z_{\mathbf{S}}^{(r=1)}(\tau, \bar{\tau}, \mathbf{z}) = \sum_{\mu \in \Lambda^* / \Lambda} Z_{\mu}(\tau) \Theta_{\mu}^*(\tau, \bar{\tau}, \mathbf{z})$$

$$Z_{\mu}(\tau) = \sum_{n=0}^{\infty} \Omega(\gamma) q^{n + \mathbf{Q}^2 / 2 - 1}, \quad \Theta_{\mu}^*(\tau, \bar{\tau}, \mathbf{z}) = \sum_{\lambda \in \mu + \Lambda} q^{-\frac{1}{2}(\lambda)_-^2} \bar{q}^{+\frac{1}{2}(\lambda)_+^2} e^{2\pi i(\lambda) \cdot z}$$

$\Omega(\gamma)$: 4d BPS index for D4-D2-D0 states (**Donaldson-Thomas invariants**)

For simplicity here focus on $\mathbf{Q} = \mathbf{Q}_- \in \Lambda_-^*$

If $\Omega(\gamma) \neq 0$ for $n = -\frac{\mathbf{Q}^2}{2} > 0$, then have states at excitation level n in 5d

Recall:

n : KK number on S_M^1 , but by level matching identified with excitation level n_L in 5d

K3 tower counting

$$\Omega(\gamma) \neq 0 \quad \text{for } \gamma = (\Sigma_{K3}, \mathbf{Q}, n) \quad \text{at } n = -\frac{\mathbf{Q}^2}{2}, \mathbf{Q} \in \Lambda_-^*$$

Key insight: [Bouchard, Creutzig, Diaconescu, Doran, Quigley, Sheshmani16]

$$\begin{aligned} Z_{\boldsymbol{\mu}}(\tau) &= \sum_{n=0}^{\infty} \Omega(\gamma) q^{n + \mathbf{Q}^2/2 - 1} \\ &= \eta^{-24}(\tau) \Phi_{\boldsymbol{\mu}}(\tau) = [q^{-1} + 24 + \mathcal{O}(q)] \Phi_{\boldsymbol{\mu}}(\tau) \end{aligned}$$

for $\Phi_{\boldsymbol{\mu}}(\tau)$ a component of a vector-valued modular form

Expansion coefficients related to **Noether-Lefschetz numbers** [Maulik, Pandharipande'13]

[Pandharipande, Thomas'16]

$$NL_{(h, \mathbf{Q})} = \text{Coeff} \left(\Phi_{\boldsymbol{\mu}}, q^{\Delta_{\text{NL}}} \right), \quad \Delta_{\text{NL}} = \frac{1}{2} \eta^{ij} Q_i Q_j + 1 - h$$

- If $\Delta_{\text{NL}} < 0$, then $NL_{(h, \mathbf{Q})} = 0$
- If $\Delta_{\text{NL}} = 0$, then $NL_{(h, \mathbf{Q})} = -2$
- If $\Delta_{\text{NL}} > 0$, then $NL_{(h, \mathbf{Q})} \in \mathbb{Z}$

\implies **States with $n = -\frac{\mathbf{Q}^2}{2}$** appear at order q^{-1} in

$$Z_{\mathbf{0}}(\tau) = \eta^{-24}(\tau) \Phi_{\mathbf{0}}(\tau) = [q^{-1} + 24 + \mathcal{O}(q)] [-2 + \mathcal{O}(q)] = -2q^{-1} + \mathcal{O}(q^0)$$

Super-extremality

$$F_{\text{Coulomb}} \stackrel{!}{\geq} F_{\text{grav}} + F_{\text{Yukawa}}$$

$$\frac{(M_{\text{Pl}} g_5^2)(Q_\alpha f^{\alpha\beta} Q_\beta)}{M_k^2 / M_{\text{Pl}}^2} \stackrel{!}{\geq} \frac{d-3}{d-2} \Big|_{d=5} + \frac{1}{2} \frac{M_{\text{Pl}}^4}{M_k^4} \left(f^{\alpha\beta} - \frac{1}{3} \hat{v}^\alpha \hat{v}^\beta \right) \partial_\alpha \left(\frac{M_k^2}{M_{\text{Pl}}^2} \right) \partial_\beta \left(\frac{M_k^2}{M_{\text{Pl}}^2} \right)$$

Explicitly check this for states at excitation level $n_k = -\frac{1}{2} \mathbf{Q}^2$

Input from string theory:

$$M_k^2 = 8\pi(n_k - 1)T_s + \Delta_{\text{CB}}$$

- First term: Contribution from string oscillators, with string tension

$$T_s = 2\pi \mathcal{V}_S M_{11d}^2 = 2\pi(4\pi)^{-2/3} \hat{\mathcal{V}}_S M_{\text{Pl}}^2$$

- Δ_{CB} : contribution from Coulomb branch in 5d

$$\Delta_{\text{CB}} = 4\pi^2 (4\pi)^{-2/3} Q_i Q_j \hat{v}^i \hat{v}^j M_{\text{Pl}}^2 \quad \hat{v}^i : \text{Kähler moduli of K3 fiber}$$

In the asymptotic limit a number of simplifications occur

\implies Together with $n_k = -\frac{1}{2} \mathbf{Q}^2$ the equality is marginally obeyed

Conclusions

Asymptotic Tower Weak Gravity Conjecture in 5d M-theory

Non-BPS tower from emergent string excitations yield WGC tower in potentially weakly coupled directions of charge lattice not associated with a KK tower

✓ Similar conclusions also in F-theory with 6d N=1 or 4d N=1

Technical question:

Improve understanding of T^4 fibrations and asymptotic Type II theories

Conceptual questions:

What if there is no (known) tower in a certain direction?

$\frac{\Lambda_{\text{WGC}}}{\Lambda_{\text{QG}}} \geq 1 \implies$ Does it makes sense to define a tower of states in the EFT?

Fact:

In many directions without a weak coupling limit BPS towers do exist

[Alim,Heidenreich,Rudelius'21] [Gendler,Heidenreich,McAllister,Moritz,Rudelius'22]

Strings and Geometry 2024 in Hamburg

